

# Iterative Reweighted Algorithms for Matrix Rank Minimization

**Karthik Mohan**

KARNA@UW.EDU

**Maryam Fazel**

MFAZEL@UW.EDU

*Department of Electrical Engineering*

*University of Washington*

*Seattle, WA 98195-4322, USA*

**Editor:** Tong Zhang

## Abstract

The problem of minimizing the rank of a matrix subject to affine constraints has applications in several areas including machine learning, and is known to be NP-hard. A tractable relaxation for this problem is nuclear norm (or trace norm) minimization, which is guaranteed to find the minimum rank matrix under suitable assumptions. In this paper, we propose a family of Iterative Reweighted Least Squares algorithms IRLS- $p$  (with  $0 \leq p \leq 1$ ), as a computationally efficient way to improve over the performance of nuclear norm minimization. The algorithms can be viewed as (locally) minimizing certain smooth approximations to the rank function. When  $p = 1$ , we give theoretical guarantees similar to those for nuclear norm minimization, that is, recovery of low-rank matrices under certain assumptions on the operator defining the constraints. For  $p < 1$ , IRLS- $p$  shows better empirical performance in terms of recovering low-rank matrices than nuclear norm minimization. We provide an efficient implementation for IRLS- $p$ , and also present a related family of algorithms, sIRLS- $p$ . These algorithms exhibit competitive run times and improved recovery when compared to existing algorithms for random instances of the matrix completion problem, as well as on the MovieLens movie recommendation data set.

**Keywords:** matrix rank minimization, matrix completion, iterative algorithms, null-space property

## 1. Introduction

The Affine Rank Minimization Problem (ARMP), or the problem of finding the minimum rank matrix in an affine set, arises in many engineering applications such as collaborative filtering (e.g., Candes and Recht, 2009; Srebro et al., 2005), low-dimensional Euclidean embedding (e.g., Fazel et al., 2003), low-order model fitting and system identification (e.g., Liu and Vandenberghe, 2008), and quantum tomography (e.g., Gross et al., 2010). The problem is as follows,

$$\begin{aligned} & \text{minimize} && \text{rank}(X) \\ & \text{subject to} && \mathcal{A}(X) = b, \end{aligned} \tag{1}$$

where  $X \in \mathbb{R}^{m \times n}$  is the optimization variable,  $\mathcal{A} : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^q$  is a linear map, and  $b \in \mathbb{R}^q$  denotes the measurements. When  $X$  is restricted to be a diagonal matrix, ARMP reduces to the cardinality

minimization or sparse vector recovery problem,

$$\begin{aligned} & \text{minimize} && \text{card}(x) \\ & \text{subject to} && Ax = b, \end{aligned}$$

where  $x \in \mathbb{R}^n$ ,  $A \in \mathbb{R}^{m \times n}$  and  $\text{card}(x)$  denotes the number of non-zeros entries of  $x$ . A commonly used convex heuristic for this problem is  $\ell_1$  minimization. The field of compressed sensing has shown that under certain conditions on the matrix  $A$ , this heuristic solves the original cardinality minimization problem (Candes and Tao, 2004). There are also geometric arguments in favor of  $\ell_1$  norm as a good convex relaxation (see, e.g., Chandrasekaran et al., 2010b, for a unifying analysis for general linear inverse problems). It has also been observed empirically (e.g., Candes et al., 2008; Lobo et al., 2006) that by appropriately weighting the  $\ell_1$  norm and iteratively updating the weights, the recovery performance of the algorithm is enhanced. It is shown theoretically that for sparse recovery from noisy measurements, this algorithm has a better recovery error than  $\ell_1$  minimization under suitable assumptions on the matrix  $A$  (Needell, 2009; Zhang, 2010). The Reweighted  $\ell_1$  algorithm has also been generalized to the recovery of low-rank matrices (Fazel et al., 2003; Mohan and Fazel, 2010a).

Another simple and computationally efficient reweighted algorithm for sparse recovery that has been proposed in the literature (Daubechies et al., 2010), is the Iterative Reweighted Least Squares algorithm (IRLS- $p$ , for any  $0 < p \leq 1$ ). Its  $k$ th iteration is given by

$$x^{k+1} = \underset{x}{\operatorname{argmin}} \left\{ \sum_i w_i^k x_i^2 : Ax = b, \right\},$$

where  $w^k \in \mathbb{R}^n$  is a weight vector with  $w_i^k = (|x_i^k|^2 + \gamma)^{p/2-1}$  ( $\gamma > 0$  being a regularization parameter added to ensure that  $w^k$  is well defined). For  $p = 1$ , a theoretical guarantee for sparse recovery has been given for IRLS- $p$  (Daubechies et al., 2010), similar to the guarantees for  $\ell_1$  minimization. The starting point of the algorithm,  $x^0$  is set to zero. So the first iteration gives the least norm solution to  $Ax = b$ . It was empirically observed (Chartrand and Staneva, 2008; Chartrand and Yin, 2008) that IRLS- $p$  shows a better recovery performance than  $\ell_1$  minimization for  $p < 1$  and has a similar performance as the reweighted  $\ell_1$  algorithm when  $p$  is set to zero. The computational benefits of IRLS-1, as well as the above mentioned theoretical and empirical improvements of IRLS- $p$ ,  $p < 1$ , over  $\ell_1$  minimization, motivate us to ask: Would an iterative reweighted algorithm bring similar benefits for recovery of low rank matrices, and would an improved performance be at the price of additional computational cost relative to the standard nuclear norm minimization?

### 1.1 Iterative Reweighted Least Squares for ARMP

Towards answering this question, we propose the iterative reweighted least squares algorithm for rank minimization, outlined below.

**Data:**  $\mathcal{A}, b$

Set  $k = 1$ . Initialize  $W_p^0 = I, \gamma^1 > 0$  ;

**while** *not converged* **do**

$X^k$	$= \underset{X}{\operatorname{argmin}} \{ \operatorname{Tr}(W_p^{k-1} X^T X) : \mathcal{A}(X) = b \}$ ;
$W_p^k$	$= (X^{kT} X^k + \gamma^k I)^{\frac{p}{2}-1}$ ;
Choose	$0 < \gamma^{k+1} \leq \gamma^k$ ;
$k$	$= k + 1$ ;

**end**

**Algorithm 1:** IRLS- $p$  Algorithm for Matrix Rank Minimization with  $0 \leq p \leq 1$

Each iteration of Algorithm 1 minimizes a weighted Frobenius norm of the matrix  $X$ , since  $\operatorname{Tr}(W_p^{k-1} X^T X) = \|(W_p^{k-1})^{1/2} X\|_F^2$ . While minimizing the Frobenius norm subject to affine constraints doesn't lead to low-rank solutions in general, through a careful reweighting of this norm we show that Algorithm 1 does indeed produce low-rank solutions under suitable assumptions. Usually a reweighted algorithm trades off computational time for improved recovery performance when compared to the unweighted convex heuristic. As an example, the reweighted  $\ell_1$  algorithm for sparse recovery (Candes et al., 2008) and the reweighted nuclear norm algorithm (Mohan and Fazel, 2010a) for matrix rank minimization solve the corresponding standard convex relaxations ( $\ell_1$  and nuclear norm minimization respectively) in their first iteration. Thus these algorithms take at least as much time as the corresponding convex algorithms. However, the iterates of the IRLS- $p$  family of algorithms simply minimize a weighted Frobenius norm, and have run-times comparable with the nuclear norm heuristic, while (for  $p < 1$ ) also enjoy improved recovery performance. In the  $p = 1$  case, we show that the algorithm minimizes a certain smooth approximation to the nuclear norm, allowing for efficient implementations while having theoretical guarantees similar to nuclear norm minimization.

## 1.2 Contributions

Contributions of this paper are as follows.

- We give a convergence analysis for IRLS- $p$  for all  $0 \leq p \leq 1$ . We also propose a related algorithm, sIRLS- $p$  (or short IRLS), which can be seen as a first-order method for locally minimizing a smooth approximation to the rank function, and give convergence results for this algorithm as well. The results exploit the fact that these algorithms can be derived from the KKT conditions for minimization problems whose objectives are suitable smooth approximations to the rank function.
- We prove that the IRLS-1 algorithm is guaranteed to recover a low-rank matrix if the linear map  $\mathcal{A}$  satisfies the following *Null space property* (NSP). We show this property is both necessary and sufficient for low-rank recovery. The specific NSP that we use first appeared in the literature recently (Oymak and Hassibi, 2010; Oymak et al., 2011), and is expressed only in terms of the singular values of the matrices in the null space of  $\mathcal{A}$ .

**Definition 1** Given  $\tau > 0$ , the linear map  $\mathcal{A} : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^p$  satisfies the  $\tau$ -Null space Property ( $\tau$ -NSP) of order  $r$  if for every  $Z \in \mathcal{N}(\mathcal{A}) \setminus \{0\}$ , we have

$$\sum_{i=1}^r \sigma_i(Z) < \tau \sum_{i=r+1}^n \sigma_i(Z) \quad (2)$$

where  $\mathcal{N}(\mathcal{A})$  denotes the null space of  $\mathcal{A}$  and  $\sigma_i(Z)$  denotes the  $i^{\text{th}}$  largest singular value of  $Z$ .

It has been shown in the literature (Oymak and Hassibi, 2010) that certain random Gaussian maps  $\mathcal{A}$  satisfy this property with high probability.

- We give a gradient projection algorithm to implement IRLS- $p$ . We extensively compare these algorithms with other state of the art algorithms on both ‘easy’ and ‘hard’ randomly picked instances of the matrix completion problem (these notions are made precise in the numerical section). We also present comparisons on the MovieLens movie recommendation data set. Numerical experiments demonstrate that IRLS-0 and sIRLS-0 applied to the matrix completion problem have a better recovery performance than the Singular Value Thresholding algorithm, an implementation of Nuclear Norm Minimization (Cai et al., 2008), on both easy and hard instances. Importantly, in the case where there is no apriori information on the rank of the low rank solution (which is common in practice), our algorithm has a significantly better recovery performance on hard problem instances as compared to other state of the art algorithms for matrix completion including IHT, FPCA (Goldfarb and Ma, 2011), and Optspace (Keshavan and Oh, 2009).

### 1.3 Related Work

We review related algorithms for recovering sparse vectors and low rank matrices. Many approaches have been proposed for recovery of sparse vectors from linear measurements including  $\ell_1$  minimization and greedy algorithms (see, e.g., Needell and Tropp, 2008; Goldfarb and Ma, 2011; Garg and Khandekar, 2009, for CoSaMP, IHT, and GraDes respectively). As mentioned earlier, reweighted algorithms including Iterative reweighted  $\ell_1$  (Candes et al., 2008) and Iterative Reweighted Least Squares (Rao and Kreutz-Delgado, 1999; Wipf and Nagarajan, 2010; Daubechies et al., 2010) with  $0 < p \leq 1$  have been proposed to improve on the recovery performance of  $\ell_1$  minimization.

For the ARMP, analogous algorithms have been proposed including nuclear norm minimization (Fazel et al., 2001), reweighted nuclear norm minimization (Mohan and Fazel, 2010a; Fazel et al., 2003), as well as greedy algorithms such as AdMiRA (Lee and Bresler, 2010) which generalizes CoSaMP, SVP (Meka et al., 2010), a hard-thresholding algorithm that we also refer to as IHT, and Optspace (Keshavan and Oh, 2009). Developing efficient implementations for nuclear norm minimization is an important research area since standard semidefinite programming solvers cannot handle large problem sizes. Towards this end, algorithms including SVT (Cai et al., 2008), NNLS (Toh and Yun, 2010), FPCA (Goldfarb and Ma, 2011) have been proposed. A spectral regularization algorithm (Mazumder et al., 2010) has also been proposed for the specific problem of matrix completion.

In a preliminary conference version of this paper (Mohan and Fazel, 2010b), we proposed the IRLS- $p$  family of algorithms for ARMP, analogous to IRLS for sparse vector recovery (Daubechies et al., 2010). The present paper gives a new and improved theoretical analysis for IRLS-1 via a simpler NSP condition, obtains complete convergence results for all  $0 \leq p \leq 1$ , and gives more extensive numerical experiments.

Independent of us and at around the same time as the publication of our conference paper, Fornasier et al. (2010) proposed the IRLSM algorithm, which is similar to our IRLS-1 algorithm but with a different weight update (a thresholding operation is used to ensure the weight matrix is invertible). The authors employ the Woodbury matrix inversion lemma to speed up the implementation of IRLSM and compare it to two other algorithms, Optspace and FPCA, showing that IRLSM has a lower relative error and comparable computational times. However, their analysis of low-rank recovery uses a Strong Null Space Property (SRNSP) (which is equivalent to the condition in, Mohan and Fazel, 2010b) and is less general than the condition we consider in this paper, as discussed in Section 3.

Also, the authors have only considered recovery of matrices of size  $500 \times 500$  in their experiments. We present extensive comparisons of our algorithms with algorithms including Optspace, FPCA, and IHT for recovery of matrices of different sizes. We observe that our IRLS and sIRLS implementations run faster than IRLSM in practice. IRLSM solves a quadratic program in each iteration using a sequence of inversions, which can be expensive for large problems, even after exploiting the matrix inversion lemma. Finally the authors of the IRLSM paper only consider the  $p = 1$  case.

The rest of the paper is organized as follows. We introduce the IRLS- $p$  algorithm for ARMP in Section 2 and give convergence and performance guarantees in Section 3. In Section 4, we discuss an implementation for IRLS- $p$ , tailored for the *matrix completion problem*, and in Section 5 we present the related algorithm sIRLS- $p$ . Numerical experiments for IRLS-0 and sIRLS-0 for the matrix completion problem, as well as comparisons with SVT, IHT, IRLSM, Optspace and FPCA are given in Section 6. The last section summarizes the paper along with future research directions.

### 1.3.1 NOTATION

Let  $\mathcal{N}(\mathcal{A})$  denote the null space of the operator  $\mathcal{A}$  and  $\text{Ran}(\mathcal{A}^*)$  denote the range space of the adjoint of  $\mathcal{A}$ . Let  $\sigma(X)$  denote the vector of decreasingly ordered singular values of  $X$  so that  $\sigma_i(X)$  denotes the  $i^{\text{th}}$  largest singular value of  $X$ . Also let  $\|X\|$ ,  $\|X\|_F$  denote the spectral norm and Frobenius norm of  $X$  respectively. The nuclear norm is defined as  $\|X\|_* = \sum_i \sigma_i(X)$ .  $I_k$  denotes the identity matrix of size  $k \times k$ .

## 2. Iterative Reweighted Least Squares (IRLS- $p$ )

In this section, we describe the IRLS- $p$  family of algorithms (Algorithm 1). Recall that replacing the rank function in (1) by  $\|X\|_*$  yields the nuclear norm heuristic,

$$\begin{aligned} & \text{minimize} && \|X\|_* \\ & \text{subject to} && \mathcal{A}(X) = b. \end{aligned}$$

We now consider other (convex and non-convex) smooth approximations to the rank function. Define the *smooth Schatten- $p$  function* as

$$\begin{aligned} f_p(X) &= \text{Tr}(X^T X + \gamma I)^{p/2} \\ &= \sum_{i=1}^n (\sigma_i^2(X) + \gamma)^{\frac{p}{2}}. \end{aligned}$$

Note that  $f_p(X)$  is differentiable for  $p > 0$  and convex for  $p \geq 1$ . With  $\gamma = 0$ ,  $f_1(X) = \|X\|_*$ , which is also known as the Schatten-1 norm. With  $\gamma = 0$  and  $p \rightarrow 0$ ,  $f_p(X) \rightarrow \text{rank}(X)$ . For  $p = 1$ , one can also derive the smooth Schatten-1 function as follows.

$$f_1(X) = \max_{Z: \|Z\|_2 \leq 1} \langle Z, X \rangle - \sqrt{\gamma} d(Z) + n\sqrt{\gamma},$$

where  $d(Z)$  is a smoothing term given by  $d(Z) = \sum_{i=1}^n (1 - \sqrt{1 - \sigma_i^2(Z)}) = n - \text{Tr}(I - Z^T Z)^{1/2}$ . Plugging in  $d(Z) = 0$  in (3) gives  $\|X\|_* + n\sqrt{\gamma}$ . Hence  $f_1(X)$  is a smooth approximation of  $\|X\|_*$  obtained by smoothing the conjugate of  $\|X\|_*$ . By virtue of the smoothing, it is easy to see that

$$\|X\|_* \leq f_1(X) \leq \|X\|_* + n\sqrt{\gamma}.$$

As  $\gamma$  approaches 0,  $f_1(X)$  becomes tightly bounded around  $\|X\|_*$ . Also let  $X^*$  be the optimal solution to minimizing  $\|X\|_*$  subject to  $\mathcal{A}(X) = b$  and let  $X^s$  be the optimal solution to minimizing  $f_1(X)$  subject to the same constraints. Then it holds that

$$(\|X\|_* - \|X^*\|_*) \leq (f_1(X) - f_1(X^s)) + n\sqrt{\gamma} \quad \forall X : \mathcal{A}(X) = b.$$

Thus, to minimize  $\|X\|_*$  to a precision of  $\epsilon$ , we need to minimize the smooth approximation  $f_1(X)$  to a precision of  $\epsilon - n\sqrt{\gamma}$ .

It is therefore of interest to consider the problem

$$\begin{aligned} &\text{minimize} && f_p(X) \\ &\text{subject to} && \mathcal{A}(X) = b, \end{aligned} \tag{3}$$

the optimality conditions of which motivate the IRLS- $p$  algorithms for  $0 \leq p \leq 1$ . We show that IRLS-1 solves the smooth Schatten-1 norm or nuclear norm minimization problem, that is, finds a globally optimal solution to (3) with  $p = 1$ . For  $p < 1$ , we show that IRLS- $p$  finds a stationary point of (3).

We now give an intuitive way to derive the IRLS- $p$  algorithm from the KKT conditions of (3). The Lagrangian corresponding to (3) is

$$L(X, \lambda) = f_p(X) + \langle \hat{\lambda}, \mathcal{A}(X) - b \rangle,$$

and the KKT conditions are  $\nabla_X L(X, \hat{\lambda}) = 0$ ,  $\mathcal{A}(X) = b$ . Note that  $\nabla f_p(X) = pX(X^T X + \gamma I)^{p/2-1}$  (see, e.g., Lewis, 1996). Letting  $\lambda = \frac{\hat{\lambda}}{p}$ , we have that the KKT conditions for (3) are given by

$$\begin{aligned} 2X(X^T X + \gamma I)^{p/2-1} + \mathcal{A}^*(\lambda) &= 0 \\ \mathcal{A}(X) &= b. \end{aligned} \tag{4}$$

Let  $W_p^k = (X^{kT}X^k + \gamma I)^{p/2-1}$ . The first condition in (4) can be written as

$$X = -\frac{1}{2}\mathcal{A}^*(\lambda)(X^T X + \gamma I)^{1-p/2}.$$

This is a fixed point equation, and a solution can be obtained by iteratively solving for  $X$  as  $X^{k+1} = \frac{1}{2}\mathcal{A}^*(\lambda)(W_p^k)^{-1}$ , along with the condition  $\mathcal{A}(X^{k+1}) = b$ . Note that  $X^{k+1}$  and the dual variable  $\lambda$  satisfy the KKT conditions for the convex optimization problem,

$$\begin{aligned} & \text{minimize} && \text{Tr } W_p^k X^T X \\ & \text{subject to} && \mathcal{A}(X) = b. \end{aligned}$$

This idea leads to the IRLS- $p$  algorithm described in Algorithm 1. Note that we also let  $p = 0$  in Algorithm 1; to derive IRLS-0, we define another non-convex surrogate function by taking limits over  $f_p(X)$ . For any positive scalar  $x$ , it holds that  $\lim_{p \rightarrow 0} \frac{1}{p}(x^p - 1) = \log x$ . Therefore,

$$\begin{aligned} \lim_{p \rightarrow 0} \frac{f_p(X) - n}{p} &= \frac{1}{2} \log \det(X^T X + \gamma I) \\ &= \sum_i \frac{1}{2} \log(\sigma_i^2(X) + \gamma). \end{aligned}$$

Thus IRLS-0 can be seen as iteratively solving (as outlined previously) the KKT conditions for the non-convex problem,

$$\begin{aligned} & \text{minimize} && \log \det(X^T X + \gamma I) \\ & \text{subject to} && \mathcal{A}(X) = b. \end{aligned} \tag{5}$$

Another way to derive the IRLS- $p$  algorithm uses an alternative characterization of the Smooth Schatten- $p$  function; see appendix A.

### 3. IRLS- $p$ : Theoretical Results

In this section, convergence properties for the IRLS- $p$  family of algorithms are studied. We also give a matrix recovery guarantee for IRLS-1, under suitable assumptions on the null space of the linear operator  $\mathcal{A}$ .

#### 3.1 Convergence of IRLS- $p$

We show that the difference between successive iterates of the IRLS- $p$  ( $0 \leq p \leq 1$ ) algorithm converges to zero and that every cluster point of the iterates is a stationary point of (3). These results generalize the convergence results given for IRLS-1 in previous literature (Mohan and Fazel, 2010b; Fornasier et al., 2010) to IRLS- $p$  with  $0 < p \leq 1$ . In this section, we drop the subscript on  $W_p^k$  for ease of notation. Our convergence analysis relies on useful auxiliary functions defined as

$$g^p(X, W, \gamma) := \begin{cases} \frac{p}{2}(\text{Tr}(W(X^T X + \gamma I)) + \frac{2-p}{p} \text{Tr}((W)^{\frac{p}{p-2}})) & \text{if } 0 < p \leq 1 \\ \text{Tr}(W(X^T X + \gamma I)) - \log \det W - n & \text{if } p = 0. \end{cases}$$

These functions can be obtained from the alternative characterization of Smooth Schatten- $p$  function with details in Appendix A. We can express the iterates of IRLS- $p$  as

$$\begin{aligned} X^{k+1} &= \operatorname{argmin}_{X: \mathcal{A}(X)=b} \mathcal{J}^p(X, W^k, \gamma^k) \\ W^{k+1} &= \operatorname{argmin}_{W>0} \mathcal{J}^p(X^{k+1}, W, \gamma^{k+1}), \end{aligned}$$

and it follows that

$$\begin{aligned} \mathcal{J}^p(X^{k+1}, W^{k+1}, \gamma^{k+1}) &\leq \mathcal{J}^p(X^{k+1}, W^k, \gamma^{k+1}) \\ &\leq \mathcal{J}^p(X^{k+1}, W^k, \gamma^k) \\ &\leq \mathcal{J}^p(X^k, W^k, \gamma^k). \end{aligned} \tag{6}$$

The following theorem shows that the difference between successive iterates converges to zero. The proof is given in Appendix B.

**Theorem 2** *Given any  $b \in \mathbb{R}^q$ , the iterates  $\{X^k\}$  of IRLS- $p$  ( $0 < p \leq 1$ ) satisfy*

$$\sum_{k=1}^{\infty} \|X^{k+1} - X^k\|_F^2 \leq 2D^{\frac{2}{p}},$$

where  $D := \mathcal{J}^p(X^1, W^0, \gamma^0)$ . In particular, we have that  $\lim_{k \rightarrow \infty} (X^k - X^{k+1}) = 0$ .

**Theorem 3** *Let  $\gamma_{\min} := \lim_{k \rightarrow \infty} \gamma^k > 0$ . Then the sequence of iterates  $\{X^k\}$  of IRLS- $p$  ( $0 \leq p \leq 1$ ) is bounded, and every cluster point of the sequence is a stationary point of (3) (when  $0 < p \leq 1$ ), or a stationary point of (5) (when  $p = 0$ ).*

**Proof** Let

$$g(X, \gamma) = \begin{cases} \operatorname{Tr}(X^T X + \gamma I)^{\frac{p}{2}} & \text{if } 0 < p \leq 1 \\ \log \det(X^T X + \gamma I) & \text{if } p = 0 \end{cases}$$

Then  $g(X^k, \gamma^k) = \mathcal{J}^p(X^k, W^k, \gamma^k)$  and it follows from (6) that  $g(X^{k+1}, \gamma^{k+1}) \leq g(X^k, \gamma^k)$  for all  $k \geq 1$ . Hence the sequence  $\{g(X^k, \gamma^k)\}$  converges. This fact together with  $\gamma_{\min} > 0$  implies that the sequence  $\{X^k\}$  is bounded.

We now show that every cluster point of  $\{X^k\}$  is a stationary point of (3). Suppose to the contrary and let  $\tilde{X}$  be a cluster point of  $\{X^k\}$  that is not a stationary point. By the definition of cluster point, there exists a subsequence  $\{X^{n_i}\}$  of  $\{X^k\}$  converging to  $\tilde{X}$ . By passing to a further subsequence if necessary, we can assume that  $\{X^{n_i+1}\}$  is also convergent and we denote its limit by  $\hat{X}$ . By definition,  $X^{n_i+1}$  is the minimizer of

$$\begin{aligned} &\text{minimize} \quad \operatorname{Tr} W^{n_i} X^T X \\ &\text{subject to} \quad \mathcal{A}(X) = b. \end{aligned} \tag{7}$$

Thus,  $X^{n_i+1}$  satisfies the KKT conditions of (7), that is,

$$X^{n_i+1} W^{n_i} \in \operatorname{Ran}(\mathcal{A}^*) \text{ and } \mathcal{A}(X^{n_i+1}) = b,$$



where  $\text{Ran}(\mathcal{A}^*)$  denotes the range space of  $\mathcal{A}^*$ . Passing to limits, we see that

$$\widehat{X}\widetilde{W} \in \text{Ran}(\mathcal{A}^*) \text{ and } \mathcal{A}(\widehat{X}) = b, \quad (8)$$

where  $\widetilde{W} = (\widetilde{X}^T \widetilde{X} + \gamma_{\min} I)^{-1}$ . From (8), we conclude that  $\widehat{X}$  is a minimizer of the following convex optimization problem,

$$\begin{aligned} & \text{minimize} && \text{Tr } \widetilde{W} X^T X \\ & \text{subject to} && \mathcal{A}(X) = b. \end{aligned} \quad (9)$$

Next, by assumption,  $\widetilde{X}$  is not a stationary point of (3) (for  $0 < p \leq 1$ ) nor (5) (for  $p = 0$ ). This implies that  $\widetilde{X}$  is not a minimizer of (9) and thus  $\text{Tr } \widetilde{W} \widehat{X}^T \widehat{X} < \text{Tr } \widetilde{W} \widetilde{X}^T \widetilde{X}$ . This is equivalent to  $\mathcal{J}^p(\widehat{X}, \widetilde{W}, \gamma_{\min}) < \mathcal{J}^p(\widetilde{X}, \widetilde{W}, \gamma_{\min})$ . From this last relation and (6) it follows that,

$$\begin{aligned} \mathcal{J}^p(\widehat{X}, \widetilde{W}, \gamma_{\min}) &< \mathcal{J}^p(\widetilde{X}, \widetilde{W}, \gamma_{\min}), \\ g(\widehat{X}, \gamma_{\min}) &< g(\widetilde{X}, \gamma_{\min}). \end{aligned} \quad (10)$$

On the other hand, since the sequence  $\{g(X^k, \gamma^k)\}$  converges, we have that

$$\lim g(X^i, \gamma^i) = \lim g(X^{n_i}, \gamma^{n_i}) = g(\widetilde{X}, \gamma_{\min}) = \lim g(X^{n_{i+1}}, \gamma^{n_{i+1}}) = g(\widehat{X}, \gamma_{\min})$$

which contradicts (10). Hence, every cluster point of  $\{X^k\}$  is a stationary point of (3) (when  $0 < p \leq 1$ ) and a stationary point of (5) (when  $p = 0$ ).  $\blacksquare$

### 3.2 Performance Guarantee for IRLS-1

In this section, we discuss necessary and sufficient conditions for low-rank recovery using IRLS-1. We show that any low-rank matrix satisfying  $\mathcal{A}(X) = b$  can be recovered via IRLS-1, if the null space of  $\mathcal{A}$  satisfies a certain property. If the desired matrix is not low-rank, we show IRLS-1 recovers it to within an error that is a constant times the best rank- $r$  approximation error, for any  $r$ . We first give a few definitions and lemmas.

**Definition 4** Given  $x \in \mathbb{R}^n$ , let  $x_{[i]}$  denote the  $i^{\text{th}}$  largest element of  $x$  so that  $x_{[1]} \geq x_{[2]} \geq \dots \geq x_{[n-1]} \geq x_{[n]}$ . A vector  $x \in \mathbb{R}^n$  is said to be majorized by  $y \in \mathbb{R}^n$  (denoted as  $x \prec y$ ) if  $\sum_{i=1}^k x_{[i]} \leq \sum_{i=1}^k y_{[i]}$  for  $k = 1, 2, \dots, n-1$  and  $\sum_{i=1}^n x_{[i]} = \sum_{i=1}^n y_{[i]}$ . A vector  $x \in \mathbb{R}^n$  is said to be weakly sub-majorized by  $y \in \mathbb{R}^n$  (denoted as  $x \prec_w y$ ) if  $\sum_{i=1}^k x_{[i]} \leq \sum_{i=1}^k y_{[i]}$  for  $k = 1, 2, \dots, n$ .

**Lemma 5** (Horn and Johnson, 1990) For any two matrices,  $A, B \in \mathbb{R}^{m \times n}$  it holds that  $|\sigma(A) - \sigma(B)| \prec_w \sigma(A - B)$ .

**Lemma 6** (Horn and Johnson, 1991; Marshall and Olkin, 1979) Let  $g : D \rightarrow \mathbb{R}$  be a convex and increasing function where  $D \subset \mathbb{R}$ . Let  $x, y \in D^n$ . Then if  $x \prec_w y$ , we have  $(g(x_1), g(x_2), \dots, g(x_n)) \prec_w (g(y_1), g(y_2), \dots, g(y_n))$ .

Since  $g(x) = (x^2 + \gamma)^{\frac{1}{2}}$  is a convex and increasing function on  $\mathbb{R}_+$ , applying Lemma 6 to the majorization inequality in Lemma 5 we have

$$\sum_{i=1}^n (|\sigma_i(A) - \sigma_i(B)|^2 + \gamma)^{\frac{1}{2}} \leq \sum_{i=1}^n (\sigma_i^2(A - B) + \gamma)^{\frac{1}{2}}. \quad (11)$$

Let  $X_0$  be a matrix (that is not necessarily low-rank) and let the measurements be given by  $b = \mathcal{A}(X_0)$ . In this section, we give necessary and sufficient conditions for recovering  $X_0$  using IRLS-1. Let  $X_0^\gamma := (X_0^T X_0 + \gamma I)^{\frac{1}{2}}$  be the  $\gamma$ -approximation of  $X_0$ , and let  $X_{0,r}, X_{0,r}^\gamma$  be the best rank- $r$  approximations of  $X_0, X_0^\gamma$  respectively.

Recall the Null space Property  $\tau$ -NSP defined earlier in (2). This condition requires that every nonzero matrix in the null space of  $\mathcal{A}$  has a rank larger than  $2r$ .

**Theorem 7** *Assume that  $\mathcal{A}$  satisfies  $\tau$ -NSP of order  $r$  for some  $0 < \tau < 1$ . For every  $X_0$  satisfying  $\mathcal{A}(X_0) = b$  it holds that*

$$f_1(X_0 + Z) > f_1(X_0), \text{ for all } Z \in \mathcal{N}(\mathcal{A}) \setminus \{0\} \text{ satisfying } \|Z\|_* \geq C \|X_0^\gamma - X_{0,r}^\gamma\|_*. \quad (12)$$

Furthermore, we have the following bounds,

$$\begin{aligned} \|\bar{X} - X_0\|_* &\leq C \|X_0^\gamma - X_{0,r}^\gamma\|_* \\ \|\bar{X}_r - X_0\|_* &\leq (2C + 1) \|X_0^\gamma - X_{0,r}^\gamma\|_*, \end{aligned}$$

where  $C = 2 \frac{(1+\tau)}{1-\tau}$ , and  $\bar{X}$  is the output of IRLS-1. Conversely, if (12) holds, then  $\mathcal{A}$  satisfies  $\delta$ -NSP of order  $r$ , where  $\delta > \tau$ .

**Proof**

Let  $Z \in \mathcal{N}(\mathcal{A}) \setminus \{0\}$  and  $\|Z\|_* \geq C \|X_0^\gamma - X_{0,r}^\gamma\|_*$ . We see that

$$\begin{aligned} f_1(X_0 + Z) &= \text{Tr}((X_0 + Z)^T (X_0 + Z) + \gamma I)^{\frac{1}{2}} = \sum_{i=1}^n (\sigma_i^2(X_0 + Z) + \gamma)^{\frac{1}{2}} \\ &\geq \sum_{i=1}^r ((\sigma_i(X_0) - \sigma_i(Z))^2 + \gamma)^{\frac{1}{2}} + \sum_{i=r+1}^n ((\sigma_i(Z) - \sigma_i(X_0))^2 + \gamma)^{\frac{1}{2}} \\ &= \sum_{i=1}^r ((\sigma_i^2(X_0) + \gamma) + \sigma_i^2(Z) - 2\sigma_i(X_0)\sigma_i(Z))^{\frac{1}{2}} \\ &\quad + \sum_{i=r+1}^n ((\sigma_i^2(Z) + \gamma) + \sigma_i^2(X_0) - 2\sigma_i(Z)\sigma_i(X_0))^{\frac{1}{2}} \\ &\geq \sum_{i=1}^r |(\sigma_i^2(X_0) + \gamma)^{\frac{1}{2}} - \sigma_i(Z)| + \sum_{i=r+1}^n |(\sigma_i^2(Z) + \gamma)^{\frac{1}{2}} - \sigma_i(X_0)| \\ &\geq \sum_{i=1}^r (\sigma_i^2(X_0) + \gamma)^{\frac{1}{2}} - \sum_{i=1}^r \sigma_i(Z) + \sum_{i=r+1}^n (\sigma_i^2(Z) + \gamma)^{\frac{1}{2}} - \sum_{i=r+1}^n \sigma_i(X_0) \\ &\geq f_1(X_0) - \sum_{i=r+1}^n (\sigma_i^2(X_0) + \gamma)^{\frac{1}{2}} - \sum_{i=1}^r \sigma_i(Z) \\ &\quad + \sum_{i=r+1}^n \sigma_i(Z) - \sum_{i=r+1}^n \sigma_i(X_0), \end{aligned} \quad (13)$$

where the first inequality follows from (11). Since  $\tau$ -NSP holds, we have

$$\begin{aligned}
 \sum_{i=1}^r \sigma_i(Z) &< \tau \sum_{i=r+1}^n \sigma_i(Z) = \sum_{i=r+1}^n \sigma_i(Z) - (1-\tau) \sum_{i=r+1}^n \sigma_i(Z) \\
 &\leq \sum_{i=r+1}^n \sigma_i(Z) - C \frac{1-\tau}{1+\tau} \|X_0^\gamma - X_{0,r}^\gamma\|_* \\
 &= \sum_{i=r+1}^n \sigma_i(Z) - 2\|X_0^\gamma - X_{0,r}^\gamma\|_*, \tag{14}
 \end{aligned}$$

where the second inequality uses  $\|Z\|_* \geq C\|X_0^\gamma - X_{0,r}^\gamma\|_*$  and  $\tau$ -NSP. Combining (13) and (14), we obtain

$$\begin{aligned}
 f_1(X_0 + Z) &= f_1(X_0) + \sum_{i=r+1}^n \sigma_i(Z) - \sum_{i=1}^r \sigma_i(Z) - \sum_{i=r+1}^n (\sigma_i^2(X_0) + \gamma)^{\frac{1}{2}} \\
 &\quad - \sum_{i=r+1}^n \sigma_i(X_0) \\
 &\geq f_1(X_0) + \sum_{i=r+1}^n \sigma_i(Z) - \sum_{i=1}^r \sigma_i(Z) - \sum_{i=r+1}^n (\sigma_i^2(X_0) + \gamma)^{\frac{1}{2}} \\
 &\quad - \sum_{i=r+1}^n (\sigma_i^2(X_0) + \gamma)^{\frac{1}{2}} \\
 &= f_1(X_0) + \sum_{i=r+1}^n \sigma_i(Z) - \sum_{i=1}^r \sigma_i(Z) - 2\|X_0^\gamma - X_{0,r}^\gamma\|_* \\
 &> f_1(X_0). \tag{15}
 \end{aligned}$$

This proves (12).

Next, if  $\widehat{X}$  is an optimal solution of (3) with  $p = 1$ , then  $Z = \widehat{X} - X_0 \in \mathcal{N}(\mathcal{A})$  and it follows immediately from (15) that

$$\|\widehat{X} - X_0\|_* = \|Z\|_* \leq C\|X_0^\gamma - X_{0,r}^\gamma\|_*.$$

Note that problem (3) is convex when  $p = 1$ , and every stationary point of (3) is a global minimum. Hence, by Theorem 3, IRLS-1 converges to a global minimum of the problem (3). It follows that  $\bar{X} = \widehat{X}$  and

$$\|\bar{X} - X_0\|_* \leq C\|X_0^\gamma - X_{0,r}^\gamma\|_*. \tag{16}$$

Finally, since  $|\sigma(\bar{X}) - \sigma(X_0)| \prec_w \sigma(\bar{X} - X_0)$ ,

$$\sum_{i=r+1}^n \sigma_i(\bar{X}) - \sum_{i=r+1}^n \sigma_i(X_0) \leq \sum_{i=1}^n |\sigma_i(\bar{X}) - \sigma_i(X_0)| \leq \|\bar{X} - X_0\|_*. \tag{17}$$

Thus,

$$\begin{aligned}
 \|\bar{X}_r - X_0\|_* &\leq \|\bar{X} - \bar{X}_r\|_* + \|\bar{X} - X_0\|_* \\
 &\leq \|X_0 - X_{0,r}\|_* + \|\bar{X} - X_0\|_* + \|\bar{X} - X_0\|_* \\
 &\leq (2C + 1)\|X_0^\gamma - X_{0,r}^\gamma\|_*,
 \end{aligned}$$

where the second inequality follows from (17) and the third inequality from (16).

Conversely, suppose that (12) holds, that is,  $f_1(X_0 + Z) > f_1(X_0)$  for all  $\|Z\|_* \geq C\|X_0^\gamma - X_{0,r}^\gamma\|_*$ ,  $Z \in \mathcal{N}(A) \setminus \{0\}$ . We would like to show that  $\mathcal{A}$  satisfies  $\delta$ -NSP of order  $r$ . Assume to the contrary that there exists  $Z \in \mathcal{N}(A)$  such that

$$\sum_{i=1}^r \sigma_i(Z) \geq \delta \sum_{i=r+1}^n \sigma_i(Z). \tag{18}$$

Let  $\alpha = \frac{1-\tau\delta}{2(1+\tau)}$  and set  $X_0 = -Z_r - \alpha(Z - Z_r)$ , where  $Z_r$  denotes the best rank- $r$  approximation to  $Z$ . Note that  $\alpha < (1 + \delta)/C$ . Assume that  $Z$  satisfies

$$\left(\frac{1+\delta}{C} - \alpha\right) \sum_{i=r+1}^n \sigma_i(Z) \geq (n-r)\sqrt{\gamma}. \tag{19}$$

If not,  $Z$  can be multiplied by a large enough positive constant so that it satisfies both (19) and (18). Note that (18) can be rewritten as,

$$\|Z\|_* \geq (1 + \delta) \sum_{i=r+1}^n \sigma_i(Z). \tag{20}$$

Combining (19) and (20), we obtain that

$$\begin{aligned} \|X_0^\gamma - X_{0,r}^\gamma\|_* &= \sum_{i=r+1}^n (\alpha^2 \sigma_i^2(Z) + \gamma)^{\frac{1}{2}} \leq (n-r)\sqrt{\gamma} + \alpha \sum_{i=r+1}^n \sigma_i(Z) \\ &\leq \frac{\|Z\|_*}{C} \end{aligned} \tag{21}$$

Moreover, it follows from (18) and the choice of  $X_0$  that

$$\begin{aligned} f_1(X_0) &= \sum_{i=1}^n (\sigma_i^2(X_0) + \gamma)^{\frac{1}{2}} \\ &\geq \sum_{i=1}^r \sigma_i(X_0) + \sum_{i=r+1}^n \sigma_i(X_0) \\ &= \sum_{i=1}^r \sigma_i(Z) + \alpha \sum_{i=r+1}^n \sigma_i(Z) \\ &\geq (\alpha + \delta) \sum_{i=r+1}^n \sigma_i(Z). \end{aligned} \tag{22}$$

On the other hand, notice by the definition of  $\alpha$  that  $\alpha > \frac{1-\delta}{2}$ . Also assume  $Z$  satisfies

$$(2\alpha + \delta - 1) \sum_{i=r+1}^n \sigma_i(Z) \geq n\sqrt{r}. \tag{23}$$

If not,  $Z$  can be multiplied by a large enough positive constant to satisfy (23) and also (18). Combining (23) and (22), we obtain further that

$$\begin{aligned} f_1(X_0 + Z) &= \sum_{i=1}^n (\sigma_i^2(X_0 + Z) + \gamma)^{\frac{1}{2}} = r\sqrt{\gamma} + \sum_{i=r+1}^n ((1-\alpha)^2 \sigma_i^2(Z) + \gamma)^{\frac{1}{2}} \\ &\leq n\sqrt{\gamma} + (1-\alpha) \sum_{i=r+1}^n \sigma_i(Z) \leq f_1(X_0). \end{aligned} \tag{24}$$

Now it is easy to see that (21) and (24) together contradicts (12), which completes the proof. ■

Thus when the sufficient condition ( $\tau$ -NSP of order  $r$ ) holds, we have shown that the best rank- $r$  approximation of the IRLS-1 solution is not far away from  $\tilde{X}$ , the solution we wish to recover, and the distance between the two is bounded by a ( $\gamma$ -approximate) rank- $r$  approximation error of  $X_0$ . It has been shown (Oymak and Hassibi, 2010) that 1-NSP of order  $r$  holds for a random Gaussian map  $\mathcal{A}$  with high probability when the number of measurements is large enough. The necessity statement can be rephrased as follows.  $\delta$ -NSP with  $\delta > \tau$  is a necessary condition for the following to hold: whenever there is an  $X$  such that  $f_1(X) \leq f_1(X_0)$ , we have that  $\|X - X_0\|_* \leq C\|X_0^\gamma - X_{0,r}^\gamma\|_*$ . Note that the necessary and sufficient conditions for recovery of approximately low-rank matrices using IRLS-1 mirror the conditions for recovery of approximately low-rank matrices using nuclear norm minimization which is that  $\mathcal{A}$  satisfy 1-NSP. We never let the weight regularization parameter  $\gamma$  go to 0, thus theoretically the solution of IRLS-1 may never exactly equal the solution of Nuclear norm minimization. However, we note that  $\gamma$  can be made very small, for example in the numerical experiments we let  $\gamma$  go to  $10^{-10}$ .

Also note that the null space property SRNSP of order  $r$ , considered in Fornasier et al. (2010) and shown to be sufficient for low-rank recovery using IRLSM, is equivalent to 1-NSP of order  $2r$ . In this paper, we show that  $\tau$ -NSP of order  $r$  (as contrasted with the order  $2r$  NSP) is both necessary and sufficient for recovery of approximately low-rank matrices using IRLS-1. Thus, our condition for the recovery of approximately low-rank matrices using IRLS-1 generalizes those stated in previous literature (Mohan and Fazel, 2010b; Fornasier et al., 2010), since it places a weaker requirement on the linear map  $\mathcal{A}$ .

#### 4. A Gradient Projection Based Implementation of IRLS

In this section, we describe IRLS-GP, a gradient projection based implementation of IRLS- $p$  for the Matrix Completion Problem.

##### 4.1 The Matrix Completion Problem

The matrix completion problem is a special case of the affine rank minimization problem with constraints that restrict some of the entries of the matrix variable  $X$  to equal given values. The problem can be written as

$$\begin{aligned} & \text{minimize} && \text{rank}(X) \\ & \text{subject to} && X_{ij} = (X_0)_{ij}, \quad (i, j) \in \Omega, \end{aligned}$$

where  $X_0$  is the matrix we would like to recover and  $\Omega$  denotes the set of entries which are revealed. We define the operator  $\mathcal{P}_\Omega : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n \times n}$  as

$$(\mathcal{P}_\Omega(X))_{ij} = \begin{cases} X_{ij} & \text{if } (i, j) \in \Omega \\ 0 & \text{otherwise.} \end{cases} \tag{25}$$

Also,  $\Omega^c$  denotes the complement of  $\Omega$ , that is, all index pairs  $(i, j)$  except those in  $\Omega$ .

## 4.2 IRLS-GP

To apply Algorithm 1 to the matrix completion problem (25), we replace the constraint  $\mathcal{A}(X) = b$  by  $\mathcal{P}_\Omega(X) = \mathcal{P}_\Omega(X_0)$ . Each iteration of IRLS solves a quadratic program (QP). A gradient projection algorithm could be used to solve the quadratic program (QP) in each iteration of IRLS. We call this implementation IRLS-GP (Algorithm 2).

**Data:**  $\Omega, \mathcal{P}_\Omega(X_0)$   
**Result:**  $\hat{X} : \mathcal{P}_\Omega(\hat{X}) = b$   
 Set  $k = 0$ . Initialize  $X^0 = 0, \gamma^0 > 0, s^0 = (\gamma^0)^{(1-\frac{p}{2})}$  ;  
**while** *IRLS iterates not converged* **do**  
      $W^k = (X^{kT} X^k + \gamma^k I)^{\frac{p}{2}-1}$ . Set  $X_{\text{temp}} = X^k$  ;  
     **while** *Gradient projection iterates not converged* **do**  
          $X_{\text{temp}} = \mathcal{P}_{\Omega^c}(X_{\text{temp}} - s^k X_{\text{temp}} W^k) + \mathcal{P}_\Omega(X_0)$ ;  
     **end**  
     Set  $X^{k+1} = X_{\text{temp}}$  ;  
     Choose  $0 < \gamma^{k+1} \leq \gamma^k, s^{k+1} = (\gamma^{k+1})^{(1-\frac{p}{2})}$  ;  
      $k = k + 1$ ;  
**end**

**Algorithm 2:** IRLS-GP for Matrix Completion

The step size used in the gradient descent step is  $s^k = 1/L^k$ , where  $L^k = 2\|W^k\|$  is the Lipschitz constant of the gradient of the quadratic objective  $\text{Tr}(W^k X^T X)$  at the  $k^{\text{th}}$  iteration. We also warm-start the gradient projection algorithm to solve for the  $(k+1)^{\text{th}}$  iterate of IRLS with the solution of the  $k^{\text{th}}$  iterate and find that this speeds up the convergence of the gradient projection algorithm in subsequent iterations. At each iteration of IRLS, computing the weighting matrix involves an inversion operation which can be expensive for large  $n$ . To work around this, we observe that the singular values of subsequent iterates of IRLS cluster into two distinct groups, so a low rank approximation of the iterates (obtained by setting the smaller set of singular values to zero) can be used to compute the weighting matrix efficiently. Computing the singular value decomposition (SVD) can be expensive. Randomized algorithms (e.g., Halko et al., 2011) can be used to compute the top  $r$  singular vectors and singular values of a matrix  $X$  efficiently, with small approximation errors, if  $\sigma_{r+1}(X)$  is small. We describe our computations of the weighting matrix below.

*Computing the weighting matrix efficiently.* Let  $U\Sigma V^T$  be the truncated SVD of  $X^k$  (keeping top  $r$  terms in the SVD with  $r$  being determined at each iteration), so that  $U \in \mathbb{R}^{m \times r}, \Sigma \in \mathbb{R}^{r \times r}, V \in \mathbb{R}^{n \times r}$ . Then for  $p = 0$ ,  $W^{k-1} = \frac{1}{\gamma^k} (U\Sigma V^T)^T (U\Sigma V^T) + \gamma^k I_n$ . It is easy to check that  $W^k = V(\gamma^k(\Sigma^2 + \gamma^k I)^{-1} - I_r)V^T + I_n$ . Thus the cost of computing the weighting matrix given the truncated SVD is  $O(nr^2)$ , saving significant computational costs. At each iteration, we choose  $r$  to be  $\min\{r_{\max}, \hat{r}\}$  where  $\hat{r}$  is the largest integer such that  $\sigma_{\hat{r}}(X^k) > 10^{-2} \times \sigma_1(X^k)$ . Also, since the singular values of  $X^k$  tend to separate into two clusters, we observe that this choice eliminates the cluster with smaller singular

values and gives a good estimate of the rank  $r$  to which  $X^k$  can be well approximated. We find that combining warm-starts for the gradient projection algorithm along with the use of randomized algorithms for SVD computations speeds up the overall computational time of the gradient projection implementation considerably.

### 5. sIRLS- $p$ : A First-order Algorithm for Minimizing the Smooth Schatten- $p$ Function

In this section, we present the sIRLS- $p$  family of algorithms that are related to the IRLS-GP implementation discussed in the previous section. We first describe the algorithm before discussing its connection to IRLS- $p$ .

**Data:**  $\mathcal{P}_\Omega, b$   
**Result:**  $\widehat{X} : \mathcal{P}_\Omega(\widehat{X}) = b$   
 Set  $k = 0$ . Initialize  $X^0 = \mathcal{P}_\Omega(X_0), \gamma^0 > 0, s^0 = (\gamma^0)^{(1-\frac{p}{2})}$  ;  
**while not converged do**  
      $W_p^k = (X^{kT} X^k + \gamma^k I)^{\frac{p}{2}-1}$  ;  
      $X^{k+1} = \mathcal{P}_{\Omega^c}(X^k - s^k X^k W_p^k) + \mathcal{P}_\Omega(X_0)$  ;  
     Choose  $0 < \gamma^{k+1} \leq \gamma^k, s^{k+1} = (\gamma^{k+1})^{(1-\frac{p}{2})}$  ;  
      $k = k + 1$   
**end**

**Algorithm 3:** sIRLS- $p$  for Matrix Completion Problem

We note that sIRLS- $p$  (Algorithm 3) is a gradient projection algorithm applied to

$$\begin{aligned} & \text{minimize} && f_p(X) = \text{Tr}(X^T X + \gamma I)^{\frac{p}{2}} \\ & \text{subject to} && \mathcal{P}_\Omega(X) = \mathcal{P}_\Omega(X_0), \end{aligned} \tag{26}$$

while sIRLS-0 is a gradient projection algorithm applied to

$$\begin{aligned} & \text{minimize} && \log \det(X^T X + \gamma I) \\ & \text{subject to} && \mathcal{P}_\Omega(X) = \mathcal{P}_\Omega(X_0). \end{aligned} \tag{27}$$

Indeed  $\nabla f_p(X^k) = X^k W_p^k$  and the gradient projection iterates,

$$\begin{aligned} X^{k+1} &= \mathcal{P}_{\{X: \mathcal{P}_\Omega(X) = \mathcal{P}_\Omega(X_0)\}}(X^k - s^k \nabla f_p(X^k)) \\ &= \mathcal{P}_{\Omega^c}(X^k - s^k \nabla f_p(X^k)) + \mathcal{P}_\Omega(X_0) \end{aligned}$$

are exactly the same as the sIRLS- $p$  (Algorithm 3) iterates when  $\gamma^k$  is a constant. In other words, for large enough  $k$  (i.e.,  $k$  such that  $\gamma^k = \gamma_{\min}$ ), the iterates of sIRLS- $p$  and sIRLS-0 are nothing but gradient projection applied to (26) and (27) respectively, with  $\gamma = \gamma_{\min}$ . We have the following convergence results for sIRLS.

**Theorem 8** *Every cluster point of sIRLS- $p$  ( $0 < p \leq 1$ ) is a stationary point of (26) with  $\gamma = \gamma_{\min}$ .*

**Theorem 9** *Every cluster point of sIRLS-0 is a stationary point of (27) with  $\gamma = \gamma_{\min}$ .*

The proof of Theorem 8 can be found in appendix C, and Theorem 9 can be proved in a similar way. With  $p = 1$ , Theorem 8 implies that sIRLS-1 has the same performance guarantee as IRLS-1 given in Theorem 7. Note that despite sIRLS- $p$  with  $p < 1$  being a gradient projection algorithm applied to non-convex problems (26) and (27), a simple step-size suffices for convergence, and we do not consider a potentially expensive line search at each iteration of the algorithm.

We now relate the sIRLS- $p$  family to the IRLS- $p$  family of algorithms. Each iteration of the IRLS- $p$  algorithm is a quadratic program, and IRLS-GP uses iterative gradient projection to solve this quadratic program. We note that sIRLS- $p$  is nothing but IRLS-GP, with each quadratic program solved approximately by terminating at the first iteration in the gradient projection inner loop. Barring this connection, the IRLS- $p$  and sIRLS- $p$  algorithms can be seen as two different approaches to solving the smooth Schatten- $p$  minimization problem (26). We examine the trade off between these two algorithms along with comparisons to other state of the art algorithms for matrix completion in the next section.

## 6. Numerical Results

In this section, we give numerical comparisons of sIRLS-0, 1 and IRLS-0, 1 with other algorithms. We begin by examining the behavior of IRLS-0 (through the IRLS-GP implementation) and its sensitivity to  $\gamma^k$  (regularization parameter in the weighting matrix,  $W^k$ ).

### 6.1 Choice of Regularization $\gamma$

We find that IRLS-0 converges faster when the regularization parameter in the weighting matrix,  $\gamma^k$ , is chosen appropriately. We consider an exponentially decreasing model  $\gamma^k = \gamma^0 / (\eta)^k$ , where  $\gamma^0$  is the initial value and  $\eta$  is a scaling parameter. We run sensitivity experiments to determine good choices of  $\gamma^0$  and  $\eta$  for the matrix completion problem. For this and subsequent experiments, the indices corresponding to the known entries  $\Omega$  are generated using i.i.d Bernoulli  $\{0, 1\}$  random variables with a mean support size  $|\Omega| = q$ , where  $q/n^2$  is the probability for an index  $(i, j)$  to belong to the support set. The completed and unknown matrix  $X_0$  of rank  $r$  is generated as  $YY^T$ , where  $Y \in \mathbb{R}^{n \times r}$  is generated using i.i.d gaussian entries. All experiments are conducted in Matlab 7.12.0 (R2011 a) on a Intel 3 Ghz core 2 duo processor with 4 GB RAM.

As will be seen from the results, the regularization parameter  $\gamma^k$  plays an important role in the recovery. We let  $\gamma^0 = \gamma_c \|X_0\|^2$  where  $\gamma_c$  is a proportional parameter that needs to be estimated. For the sensitivity analysis of IRLS-0 (with respect to  $\gamma^0$  and  $\eta$ ), we consider matrices of size  $500 \times 500$ .

#### 6.1.1 CHOICE OF $\gamma_c$

As can be seen from Figures 1 a) and b), choosing  $\gamma_c$  appropriately leads to a better convergence rate for IRLS-0. Small values of  $\gamma_c$  ( $< 10^{-3}$ ) don't give good recovery results (premature convergence to a larger relative error). However larger values of  $\gamma_c$  ( $> 1$ ) might lead to a delayed convergence. As a heuristic, we observe that  $\gamma_c = 10^{-2}$  works well. We also note that this choice of  $\gamma_c$  works well



even if the spectral norm of  $X_0$  varies from 1 to 1000. Thus for future experiments, we normalize  $X_0$  to have a spectral norm of 1.

### 6.1.2 CHOICE OF $\eta$

Figures 2 a),b),c), and d) look at the sensitivity of the IRLS-0 algorithm to the scaling parameter,  $\eta$ . We observe that for a good choice of  $\gamma^0$  (described earlier),  $\eta$  depends on the rank of  $X_0$  to be recovered. More specifically,  $\eta$  seems to have an inverse relationship with the rank of  $X_0$ . From Figures 2 a) and d), it is clear that  $\eta = 1.3$  works well if rank of  $X_0$  equals 2 and  $\eta = 1.05$  works well when rank of  $X_0$  equals 15. More generally, the choice of  $\eta$  seems to depend on the hardness of the problem instance being considered. We formalize this notion in the next section.




Figure 1:  $n = 500$ ,  $r = 5$ ,  $\eta = 1.15$ .  $\gamma^0 = \gamma_c \|X_0\|^2$ . From left to right: Recovery error using IRLS-0 for  $\|X_0\| = 1$ ,  $\|X_0\| = 1000$ .

## 6.2 Numerical Experiments

We classify our numerical experiments into two categories based on the *degrees of freedom ratio*, given by  $FR = r(2n - r)/q$ . Note that for a  $n \times n$  matrix of rank  $r$ ,  $r(2n - r)$  is the number of degrees of freedom in the matrix. Thus if  $FR$  is large (close to 1), recovering  $X_0$  becomes harder (as the number of measurements is close to the degrees of freedom) and conversely if  $FR$  is close to zero, recovering  $X_0$  becomes easier.

We conduct subsequent numerical experiments over what we refer to in this paper as *Easy problems* ( $FR < 0.4$ ) and *Hard problems* ( $FR > 0.4$ ). We define the recovery to be successful when the relative error,  $\|X - X_0\|_F / \|X_0\|_F \leq 10^{-3}$  (with  $X$  being the output of the algorithm considered) and unsuccessful recovery otherwise. For each problem (easy or hard), the results are reported over 10 random generations of the support set,  $\Omega$  and  $X_0$ . We use NS to denote the number of successful recoveries for a given problem. Also, computation times are reported in seconds. For sIRLS and IRLS-GP (implementation of IRLS), we fix  $\eta = 1.1$  if  $FR < 0.4$  and  $\eta = 1.03$  if  $FR > 0.4$  based




Figure 2:  $n = 500$ ,  $\gamma_c = 10^{-2}$ . Clockwise from top left: Recovery error using IRLS-0 for ranks 2, 5, 10, 15 respectively.

on our earlier observations. In the next few sections, we compare the IRLS implementations with other state of the art algorithms on both exact and noisy matrix completion problems.

### 6.3 Comparison of (s)IRLS and Nuclear Norm Minimization

In this section, we compare the gradient projection implementation IRLS-GP (of IRLS-0,1) and the algorithm sIRLS-0,1 with the *Singular Value Thresholding* (SVT) algorithm (an implementation for nuclear norm minimization Cai et al., 2008) on both easy and hard problem sets. Note that SVT is not the only implementation of nuclear norm minimization. Other implementations include NNLS (Toh and Yun, 2010) and Spectral Regularization (Mazumder et al., 2010).

When we refer to IRLS-0,1 in the tables and in subsequent paragraphs, we mean their gradient projection implementation, IRLS-GP. We compare (s)IRLS-0,1 and SVT in Tables 1 and 3. A few aspects of these comparisons are highlighted below.

Problem				IRLS-1		sIRLS-1		IRLS-0	
$n$	$r$	$\frac{q}{n^2}$	$FR$	# iter	Time	# iter	Time	# iter	Time
100	10	0.57	0.34	133	4.49	132	1.63	54	0.79
200	10	0.39	0.25	140	4.49	140	2.41	60	1.34
500	10	0.2	0.2	160	24.46	163	8	77	9.63
500	10	0.12	0.33	271	37.47	336	13.86	220	22.74
1000	10	0.12	0.17	180	113.72	195	32.21	109	55.42
1000	50	0.39	0.25	140	134.30	140	102.64	51	59.74
1000	20	0.12	0.33	241	156.09	284	57.85	188	96.20
2000	20	0.12	0.17	180	485.24	190	166.28	100	235.94
2000	40	0.12	0.33	236	810.13	270	322.96	170	432.34

Table 1: Comparison of IRLS-0,1 and sIRLS-1. Performance on Easy Problems  $FR < 0.4$ .

Problem				sIRLS-1		sIRLS-0		SVT	
$n$	$r$	$\frac{q}{n^2}$	$FR$	# iter	Time	# iter	Time	# iter	Time
100	10	0.57	0.34	132	1.63	59	0.84	170	5.69
200	10	0.39	0.25	140	2.41	63	1.31	109	3.74
500	10	0.2	0.2	163	8	98	4.97	95	5.9
500	10	0.12	0.33	336	13.86	280	11.03	-	-
1000	10	0.12	0.17	195	32.21	140	20.80	85	10.71
1000	50	0.39	0.25	140	102.64	60	61.32	81	49.17
1000	20	0.12	0.33	284	57.85	241	43.11	-	-
2000	20	0.12	0.17	190	166.28	130	98.55	73	42.31
2000	40	0.12	0.33	270	322.96	220	227.07	-	-

Table 2: Comparison of sIRLS-0,1 with SVT. Performance on Easy Problems  $FR < 0.4$ .

Problem				sIRLS-1			IRLS-0			sIRLS-0		
$n$	$r$	$\frac{q}{n^2}$	$FR$	# iter	NS	Time	# iter	NS	Time	# iter	NS	Time
40	9	0.5	0.8	4705	4	163.2	1385	10	17.36	2364	9	30.22
100	14	0.3	0.87	10000	0	545.91	4811	10	89.51	5039	7	114.54
500	20	0.1	0.78	10000	0	723.58	4646	8	389.66	5140	10	315.57
1000	20	0.1	0.4	645	10	142.84	340	10	182.78	406	10	97.15
1000	20	0.06	0.66	10000	0	1830.98	2679	10	921.15	2925	10	484.84
1000	30	0.1	0.59	1152	10	295.56	781	10	401.98	915	10	244.23
1000	50	0.2	0.49	550	10	342	191	10	239.77	270	10	234.25

Table 3: Comparison of sIRLS-1, IRLS-0 and sIRLS-0. Performance on Hard Problems  $FR \geq 0.4$

### 6.3.1 IRLS-0 vs IRLS-1

Between IRLS-0 and IRLS-1, IRLS-0 takes fewer iterations to converge successfully and has a lower computational time (Table 1). The same holds true between sIRLS-0 and sIRLS-1. sIRLS-0 is also successful on more hard problem instances than sIRLS-1 (Table 3). This indicates that (s)IRLS- $p$  with  $p = 0$  has a better recovery performance and computational time as compared to  $p = 1$ .

### 6.3.2 IRLS vs sIRLS

Between sIRLS and IRLS, sIRLS-1 takes more iterations to converge as compared to IRLS-1. However because it has a lower per iteration cost, sIRLS-1 takes significantly lower computational time than IRLS-1 (Table 1). The same holds true for sIRLS-0. Thus sIRLS-0,1 are not only simpler algorithms, they also have a lower overall run time as compared to IRLS-0,1.

### 6.3.3 COMPARISON ON EASY PROBLEMS

Table 2 shows that sIRLS-0 and sIRLS-1 have competitive computational times as compared to SVT (implementation available at, Candes and Becker, 2010). There are also certain instances where SVT fails to have successful recovery while sIRLS-1 succeeds. Thus sIRLS-1 is competitive and in some instances better than SVT.

### 6.3.4 COMPARISON ON HARD PROBLEMS

For hard problems, Table 3 shows that sIRLS-0 and IRLS-0 are successful in almost all problems considered, while sIRLS-1 is not successful in 4 problems. We also found that SVT was not successful in recovery for any of the hard problems. (s)IRLS-0 also compares favorably with FPCA (Goldfarb and Ma, 2011) and Optspace (Keshavan and Oh, 2009) in terms of recovery and computational time on the easy and hard problem sets. These results are given subsequently.

In summary, (s)IRLS-1,(s)IRLS-0 have a better recovery performance than a nuclear norm minimization implementation (SVT) as evidenced by successful recovery in both easy and hard problem sets. We note that (s)IRLS-1 converges to the Nuclear Norm Minimizer (when the regularization,  $\gamma \rightarrow 0$ ) and empirically has a better recovery performance than SVT. We also note that among the family of (s)IRLS- $p$  algorithms tested, sIRLS-0 and IRLS-0 are better in both recovery performance and computational times.

## 6.4 Comparison of Algorithms for Exact Matrix Completion

As observed in the previous section, sIRLS has a lower total run time compared to IRLS-GP. Thus in subsequent experiments we compare other algorithms only with sIRLS.

### 6.4.1 DESIGN OF EXPERIMENTS

In this section, we report results from two sets of experiments. In the first set, we compare sIRLS-0 (henceforth referred to as sIRLS), *Iterative Hard Thresholding* algorithm (IHT) (Goldfarb and

Problem				sIRLS		IRLSM		IHT		Optspace	
$n$	$r$	$\frac{q}{n^2}$	$FR$	# iter	Time	#iter	Time	#iter	Time	#iter	Time
100	10	0.57	0.34	56	0.8	68	0.58	44	0.51	27	0.6
200	10	0.39	0.25	61	0.96	78	1.53	53	0.95	19	1.28
500	10	0.2	0.2	99	4.5	106	15	105	3.63	18	8.08
500	10	0.12	0.33	285	13.24	240	160.62	344	12.50	29	12.45
1000	10	0.12	0.17	143	21.17	152	106.18	192	19.30	16	28.93
1000	50	0.39	0.25	60	27.39	60	429.25	46	19.58	17	1755.44
1000	20	0.12	0.33	244	45.33	264	396.49	289	40.39	38	241.94
2000	20	0.12	0.17	130	82.47	140	916.48	179	80.76	14	428.37
2000	40	0.12	0.33	230	229.53	220	4213.4	270	225.46	28	4513

Table 4: Comparison of sIRLS-0, IRLSM, IHT and Optspace on Easy Problems with rank of the matrix to be recovered known apriori.

Ma, 2011; Meka et al., 2010), Optspace and IRLSM (Fornasier et al., 2010) over easy and hard problems with the assumption that the rank of the matrix to be recovered ( $X_0$ ) is known. We use the implementations of IRLSM, Optspace available on the authors webpage (Fornasier et al., 2012; Keshavan et al., 2009a). When the rank of  $X_0$  (denoted as  $r$ ) is known, the weighting matrix  $W^k$  for sIRLS is computed using a rank  $r$  approximation of  $X^k$  (also see Section 3.1). The second set of experiments correspond to the case where the rank of  $X_0$  is unknown, which is a more practical assumption.

#### 6.4.2 RANK OF $X_0$ KNOWN APRIORI

All the algorithms are fully successful (NS = 10) on the easy problem sets. As seen in Table 4, Optspace takes fewer iterations to converge as compared to sIRLS, IRLSM and IHT. On the other hand, sIRLS and IHT are significantly faster than Optspace and much faster than IRLSM (Fornasier et al., 2010). Although IRLSM uses a sub-routine implemented in C to speed up inverse matrix computations inside a Matlab code, it is still about 10 times slower than sIRLS on easy problems and even slower on hard problems. We note that for the hard problems in Table 5, the rank of the true solution is given by (9,14,20,20,20,30,50) respectively. IRLSM is also not successful on any of the instances for two of the hard problems. Also while sIRLS, Optspace and IHT are fully successful on most of the hard problems (see Table 5), Optspace takes considerably higher time as compared to IHT and sIRLS. Thus, when the rank of  $X_0$  is known, sIRLS is competitive with IHT in performance and computational time and much faster than Optspace and IRLSM.

#### 6.4.3 RANK OF $X_0$ UNKNOWN

A possible disadvantage of IHT and Optspace could be their sensitivity to the knowledge of the rank of  $X_0$ . Thus, our second set of experiments compare sIRLS, IHT, Optspace and FPCA (Goldfarb and Ma, 2011) over easy and hard problems when the rank of  $X_0$  is unknown. We use a heuristic for

Problem		sIRLS-0			IRLSM			IHT			Optspace		
$n$	$\frac{q}{n^2}$	# iter	NS	Time	# iter	NS	Time	# iter	NS	Time	# iter	NS	Time
40	0.5	1718	10	12.67	-	0	-	1635	10	12.16	1543	7	6.82
100	0.3	4298	8	60.18	-	0	-	4868	10	68.56	4011	5	131.62
1000	0.1	417	10	78.26	428	10	642.53	466	10	65.84	69	10	409.66
1000	0.08	814	10	151.86	1120	10	1674.2	947	10	134.30	103	10	580.09
1000	0.07	1368	10	251.46	2180	8	3399.2	1564	10	225.13	147	10	806.32
1000	0.1	949	10	226.12	1536	10	4451.1	1006	10	189.33	134	10	1904.47
1000	0.2	270	10	123.84	244	10	1736	254	10	105.88	46	10	2968

Table 5: Comparison of sIRLS-0, IHT and Optspace on Hard Problems with rank of the matrix to be recovered known apriori.

Problem				sIRLS		IHT		FPCA	Optspace	
$n$	$r$	$\frac{q}{n^2}$	$FR$	# iter	Time	# iter	Time	Time	# iter	Time
100	10	0.57	0.34	59	1.61	38	1.15	0.13	25	0.62
200	10	0.39	0.25	62	2.55	44	1.96	0.37	17	1.4
500	10	0.2	0.2	98	9.39	71	8.65	2.52	17	8.46
500	10	0.12	0.33	283	16.07	225	23.23	71.26	30	11.63
1000	10	0.12	0.17	140	38.44	104	31.47	11.24	14	26.31
1000	50	0.39	0.25	60	217.79	35	132.36	15.10	17	1774.08
1000	20	0.12	0.33	241	77.52	177	70.13	18.51	30	199.98
2000	20	0.12	0.17	130	236.19	98	152.36	42.06	12	374.15
2000	40	0.12	0.33	220	234.44	167	323.67	76.26	26	3466

Table 6: Comparison of sIRLS-0, IHT, FPCA and Optspace on Easy Problems when no prior information is available on the rank of the matrix to be recovered.

determining the approximate rank of  $X^k$  at each iteration for sIRLS, IHT. Heuristics for determining approximate rank are also used in the respective implementations for FPCA and Optspace. We note that computing the approximate rank is important for speeding up the SVD computations in all of these algorithms.

*Choice of rank.* We choose  $r$  (the rank at which the SVD of  $X^k$  is truncated) to be  $\min\{r_{\max}, \hat{r}\}$  where  $\hat{r}$  is the largest integer such that  $\sigma_{\hat{r}}(X^k) > \alpha \times \sigma_1(X^k)$ . For IHT we find that  $\alpha = 5 \times 10^{-2}$  works well while for sIRLS and FPCA,  $\alpha = 10^{-2}$  works well. The SVD computations in IHT, sIRLS are based on a randomized algorithm (Halko et al., 2011). We note that Linear-Time SVD (Drineas et al., 2006) is used to compute the SVD in the FPCA implementation (Goldfarb and Ma, 2009), and although faster than the randomized SVD algorithm we use for sIRLS, it can be significantly less accurate.

Problem		sIRLS			IHT			FPCA		Optspace		
$n$	$FR$	# iter	NS	Time	# iter	NS	Time	NS	Time	# iter	NS	Time
40	0.8	1498	10	12.91	-	0	-	5	1.69	-	0	-
100	0.87	4934	5	72.36	-	0	-	0	-	-	0	-
500	0.78	4859	9	326.06	-	0	-	0	-	-	0	-
1000	0.40	406	10	115.73	280	10	72.67	10	26.54	40	10	256.92
1000	0.57	1368	10	237.22	1059	10	244.49	0	-	133	5	769.29
1000	0.66	2961	10	554.25	-	0	-	0	-	-	0	-
1000	0.59	897	10	276.08	660	10	213.95	10	62.43	89	5	1420.81
1000	0.49	270	10	263.45	203	10	186.15	10	25.21	45	10	2924.68

Table 7: Comparison of sIRLS-0, IHT, FPCA and Optspace on Hard Problems when no prior information is available on the rank of the matrix to be recovered.

*Comparison of algorithms.* All the algorithms compared are successful on the easy problems. However, Optspace takes much more time to converge on recovering matrices with high rank as can be seen from Table 6. sIRLS, FPCA and IHT have competitive run times on all the problems. For hard problems, however, sIRLS has a clear advantage over IHT, Optspace and FPCA in successful recovery (Table 7). sIRLS is fully successful on all problems except the second and third on which it has a success rate of 5 and 9 respectively. On the other hand IHT, Optspace and FPCA have partial or unsuccessful recovery in many problems. sIRLS is competitive with IHT and FPCA on computational times while Optspace is much slower than all the other algorithms. Thus, when the rank of  $X_0$  is not known apriori, sIRLS has a distinct advantage over IHT, Optspace and FPCA in successfully recovering  $X_0$  for hard problems.

## 6.5 Comparison of Algorithms for Noisy Matrix Completion

In this subsection, we compare IHT, FPCA and sIRLS on randomly generated noisy matrix completion problems. We consider the following noisy matrix completion problem,

$$\begin{aligned} & \text{minimize} && \text{rank}(X) \\ & \text{subject to} && \mathcal{P}_\Omega(X) = \mathcal{P}_\Omega(B), \end{aligned}$$

where  $\mathcal{P}_\Omega(B) = \mathcal{P}_\Omega(X_0) + \mathcal{P}_\Omega(Z)$ ,  $X_0$  is a low rank matrix of rank  $r$  that we wish to recover and  $\mathcal{P}_\Omega(Z)$  is the measurement noise. Note that this noise model has been used before for matrix completion (see Cai et al., 2008). Let  $Z_{ij}$  be i.i.d Gaussian random variables with distribution  $\mathcal{N}(0, \sigma^2)$ . We would like the noise to be such that  $\|\mathcal{P}_\Omega(Z)\|_F \leq \epsilon \|\mathcal{P}_\Omega(X_0)\|_F$  for a noise parameter  $\epsilon$ . This would be true if  $\sigma \sim \epsilon\sqrt{r}$  (Cai et al., 2008).

We adapt sIRLS for noisy matrix completion by replacing  $\mathcal{P}_\Omega(X_0)$  by  $\mathcal{P}_\Omega(B)$  in Algorithm 3. For all the algorithms tested in Table 8, we declare the recovery to be successful if  $\|X - X_0\|_F / \|X_0\|_F \leq \epsilon = 10^{-3}$ , where  $X$  is the output of the algorithms. Table 8 shows that sIRLS has successful recovery for easy noisy matrix completion problems with apriori knowledge of rank. The same holds true for

Problem				sIRLS			IHT			Optspace		
$n$	$r$	$\frac{q}{n^2}$	$FR$	# iter	NS	Time	#iter	NS	Time	#iter	NS	Time
100	10	0.57	0.34	51	10	0.55	41	10	0.44	28	10	0.29
200	10	0.39	0.25	56	10	0.78	48	10	0.60	19	10	0.56
500	10	0.2	0.2	96	10	4.11	88	10	2.76	18	10	6.82
500	10	0.12	0.33	298	10	12.60	298	10	9.56	29	10	12.45
1000	10	0.12	0.17	141	10	19.66	132	10	12.28	15	10	37.17
1000	10	0.39	0.25	50	10	20.94	40	10	15.53	18	10	1197.48
1000	20	0.12	0.33	254	10	44.35	247	10	32.03	25	10	220.60
2000	20	0.12	0.17	130	10	77.30	121	10	84.23	12	10	469.67
2000	40	0.12	0.33	236	10	221.31	227	10	170.88	28	10	4515.06

Table 8: Comparison of sIRLS, IHT and Optspace on the Noisy Matrix Completion problem.

hard problems with the true rank known apriori. Thus sIRLS has a competitive performance even for noisy recovery.

## 6.6 Application to Movie Lens Data Set

In this section, we consider the movie lens data sets (Dahlen et al, 1998) with 100,000 ratings. In particular, we consider four different splits of the 100k ratings into (training set, test set): (u1.base, u1.test), (u2.base, u2.test), (u3.base, u3.test), (u4.base, u4.test) for our numerical experiments. Any given set of ratings (e.g., from a data split) can be represented as a matrix. This matrix has rows representing the users and columns representing the movies and an entry  $(i, j)$  of the matrix is non-zero if we know the rating of user  $i$  for movie  $j$ . Thus estimating the remaining ratings in the matrix corresponds to a matrix completion problem. For each data split, we train sIRLS, IHT, and Optspace on the training set and compare their performance on the corresponding test set. The performance metric here is *Normalized Mean Absolute Error* or NMAE given as follows. Let  $M$  be the matrix representation corresponding to the actual test ratings and  $X$  be the ratings matrix output by an algorithm when input the training set. Then

$$\text{NMAE} = \left( \sum_{i,j \in \text{supp}(M)} \frac{|M_{ij} - X_{ij}|}{|\text{supp}(M)|} \right) / (rt_{\max} - rt_{\min}),$$

where  $rt_{\min}$  and  $rt_{\max}$  are the minimum and maximum movie ratings possible. The choice of  $\gamma, \eta$  for sIRLS is the same as for the random experiments (described in previous sections). sIRLS is terminated if the maximum number of iterations exceeds 700 or if the relative error between the successive iterates is less than  $10^{-3}$ . We set the rank of the unknown ratings matrix to be equal to 5 while running all the three algorithms. Table 9 shows that the NMAE for sIRLS, IHT, and Optspace are almost the same across different splits of the data. Keshavan et al. (2009b) reported the NMAE results for different algorithms when tested on data split 1. These were reported to be 0.186, 0.19 and 0.242 for Optspace (Keshavan and Oh, 2009), FPCA (Goldfarb and Ma, 2011), and AdMiRA



	sIRLS	IHT	Optspace
split 1	0.1919	0.1925	0.1887
split 2	0.1878	0.1883	0.1878
split 3	0.1870	0.1872	0.1881
split 4	0.1899	0.1896	0.1882

Table 9: NMAE for sIRLS-0 for different splits of the 100k movie-lens data set.

(Lee and Bresler, 2010) respectively. Thus sIRLS has a NMAE that is as good as Optspace, FPCA, IHT and has a better NMAE than AdMiRA.

## 7. Conclusions and Future Directions

We proposed a family of Iterative Reweighted Least Squares algorithms (IRLS- $p$ ) for the affine rank minimization problem. We showed that IRLS-1 converges to the global minimum of the smoothed nuclear norm, and that IRLS- $p$  with  $p < 1$  converges to a stationary point of the corresponding non-convex yet smooth approximation to the rank function. We gave a matrix recovery guarantee for IRLS-1, showing that it approximately recovers the true low-rank solution (within a small error depending on the algorithm parameter  $\gamma$ ), if the null space of the affine measurement map satisfies certain properties. This null space condition is both necessary and sufficient for low-rank recovery, thus improving on and simplifying the previous analysis for IRLS-1 (Mohan and Fazel, 2010b).

We then focused on the matrix completion problem, a special case of affine rank minimization arising in collaborative filtering among other applications, and presented efficient implementations specialized to this problem. We gave an implementation for IRLP- $p$  for this problem using gradient projections. We also presented a related first-order algorithm, sIRLS- $p$ , for minimizing the smooth Schatten- $p$  function, which serves as a smooth approximation of the rank. Our first set of numerical experiments show that (s)IRLS-0 has a better recovery performance than nuclear norm minimization via SVT. We show that sIRLS-0 has a good recovery performance even when noise is present. Our second set of experiments demonstrate that sIRLS-0 compares favorably in terms of performance and run time with IHT, Optspace, and IRLSM when the rank of the low rank matrix to be recovered is known. When the rank information is absent, sIRLS-0 shows a distinct advantage in performance over IHT, Optspace and FPCA.

### 7.1 Future Directions

Low-rank recovery problems have recently been pursued in machine learning motivated by applications including collaborative filtering. Iterative reweighted algorithms for low-rank matrix recovery have empirically exhibited improved performance compared to unweighted convex relaxations. However, there has been a relative lack of theoretical results, as well as efficient implementations for these algorithms. This paper takes a step in addressing both of these issues, and opens up several directions for future research.

*Low-rank plus sparse decomposition.* The problem of decomposing a matrix into a low-rank component and a sparse component has received much attention (Chandrasekaran et al., 2011; Tan et al., 2011), and arises in graphical model identification (Chandrasekaran et al., 2010a) as well as a version of robust PCA (Candes et al., 2011), where problem sizes of practical interest are often very large. The convex relaxation proposed for this problem minimizes a combination of nuclear norm and  $\ell_1$  norm. An interesting direction for future work is to extend the IRLS algorithms family to this problem, by combining the vector and the matrix weighted updates. A potential feature of such an algorithm can be that the value of  $p$  for the vector part and the matrix part (and hence the weights) can be chosen separately, allowing control over how aggressively to promote the sparsity and the low-rank features.

*Distributed IRLS.* In the IRLS family, the least squares problem that is solved in every iteration is in fact *separable* in the columns of the matrix  $X$  (as also pointed out in Fornasier et al., 2010), so it can be solved completely in parallel. This opens the door not just to a fast parallel implementation, but also to the possibility of a partially distributed algorithm. Noting that the weight update step does not appear easy to decompose, an interesting question is whether we can use approximate but decomposable weights, so that the updates would require only local information.

*Other applications for the NSP.* The simple Null space Property used here, being based on only the singular values of elements in the null space, makes the connection between associated vector and matrix recovery proofs clear and transparent, and may be of independent interest (see Oymak et al., 2011).

## Acknowledgments

We would like to acknowledge Ting Kei Pong for helpful discussions on the algorithms and Samet Oymak for pointing out an important singular value inequality. M. F. also acknowledges helpful discussions with Holger Rauhut.

## Appendix A. IRLS- $p$ from Characterization of Smooth Schatten- $p$ Function

We can also derive the IRLS- $p$  algorithm for  $0 < p \leq 1$  by defining the following function,

$$\mathcal{F}^p(X, W, \gamma) = \text{Tr} W^{\frac{p-2}{p}} (X^T X + \gamma I)$$

The following lemma (Argyriou et al., 2007; Argyriou, 2010) proves useful for our analysis.

**Lemma 10** *Let  $\gamma > 0$  and*

$$W^* = \frac{(X^T X + \gamma I)^{\frac{p}{2}}}{\text{Tr}(X^T X + \gamma I)^{\frac{p}{2}}}.$$

*Then*

$$W^* = \underset{W}{\text{argmin}} \{ \mathcal{F}^p(X, W, \gamma) : W \succ 0, \text{Tr}(W) \leq 1 \}.$$

Note that  $\mathcal{F}^p(X, W^*, \gamma) = (f_p(X))^{\frac{2}{p}}$ . Hence the problem of minimizing the *Smooth Schatten- $p$  function*  $f_p(X)$  (3) is equivalent to the following

$$\begin{aligned} & \text{minimize} && \mathcal{F}^p(X, W, \gamma) \\ & \text{subject to} && \mathcal{A}(X) = b, W \succ 0, \text{Tr} W \leq 1, \end{aligned} \quad (28)$$

where the variables are  $X$  and  $W$ . As a relaxation to minimizing (28) jointly in  $X, W$ , one can consider minimizing (28) alternately with respect to  $X$  and  $W$  as in Algorithm 4.

**Data:**  $\mathcal{A}, b$   
**Result:**  $\hat{X} : \mathcal{A}(\hat{X}) = b$   
 Set  $k = 0$ . Initialize  $W_p^0 = I, \gamma^1 > 0$  ;  
**while not converged do**  
      $X^{k+1} = \underset{X}{\text{argmin}} \mathcal{F}^p(X, W^k)$  ;  
         s.t.  $\mathcal{A}(X) = b$  ;  
      $W^{k+1} = \underset{W}{\text{argmin}} \mathcal{F}^p(X^{k+1}, W)$  ;  
         s.t.  $W \succ 0, \text{Tr}(W) \leq 1$  ;  
     Choose  $0 < \gamma^{k+2} \leq \gamma^{k+1}$  ;  
      $k = k + 1$  ;  
**end**

**Algorithm 4:** Alternative representation of the IRLS- $p$  algorithm.

We note (and shall prove) that Algorithm 4 gives us the same  $X$  updates as the IRLS- $p$  algorithm. This gives an interpretation to the IRLS- $p$  algorithm as alternately minimizing an equivalent *Smooth Schatten- $p$  problem* (28). Consider minimization with respect to  $W$  of  $\mathcal{F}^p(X, W, \gamma)$  with  $X$  fixed. This problem can be re-formulated as

$$\begin{aligned} & \text{minimize} && \text{Tr} \hat{W} (X^T X + \gamma I) \\ & \text{subject to} && \text{Tr} \hat{W}^{\frac{p}{p-2}} \leq 1 \\ & && \hat{W} \succ 0, \end{aligned} \quad (29)$$

where  $\hat{W} = W^{\frac{p-2}{p}}$ . The following lemma relates  $\mathcal{F}^p(X, W, \gamma)$  with  $\mathcal{J}^p(X, W, \gamma)$  (defined in (6)).

**Lemma 11** *Let*

$$\hat{\mathcal{W}}^* = \frac{(X^T X + \gamma I)^{\frac{p}{2}-1}}{(\text{Tr}(X^T X + \gamma I)^{\frac{p}{2}})^{\frac{p-2}{p}}}.$$

*Then  $\hat{\mathcal{W}}^*$  is the optimal solution to (29) as well as the following problem:*

$$\begin{aligned} & \text{minimize} && \text{Tr} \hat{W} (X^T X + \gamma I) + \lambda (\text{Tr} \hat{W}^{\frac{p}{p-2}} - 1), \\ & \text{subject to} && \hat{W} \succ 0, \end{aligned}$$

where  $\lambda = \frac{2-p}{p} (\text{Tr}(X^T X + \gamma I)^{\frac{p}{2}})^{\frac{2}{p}}$ . Furthermore, let

$$\tilde{W} = \underset{W \succ 0}{\text{argmin}} \mathcal{J}^p(X, W, \gamma).$$

Then  $\mathcal{J}^p(X, \tilde{W}, \gamma) = f_p(X)$  and

$$\operatorname{argmin}_X \left\{ \mathcal{F}^p(X, (\widehat{\mathcal{W}}^*)^{\frac{p}{p-2}}, \gamma) : \mathcal{A}(X) = b \right\} = \operatorname{argmin}_Z \left\{ \mathcal{J}^p(Z, \tilde{W}, \gamma) : \mathcal{A}(Z) = b \right\}.$$

Thus Lemma (11) shows that alternately minimizing  $\mathcal{F}^p(X, W, \gamma)$  with respect to  $W$  followed by  $X$  (with constraints on  $W$  as in (29) and affine constraints on  $X$ ) is equivalent to alternately minimizing  $\mathcal{J}^p(X, W, \gamma)$  with respect to  $W$  followed by  $X$  (with affine constraints on  $X$  and  $\{W : W \succ 0\}$ ).

**Proof** The Lagrangian for (29) is given by

$$L(\widehat{W}, \lambda) = \operatorname{Tr} \widehat{W} (X^T X + \gamma I) + \lambda (\operatorname{Tr} \widehat{W}^{\frac{p}{p-2}} - 1)$$

Note that  $\lambda^* = \frac{2-p}{p} (\operatorname{Tr}(X^T X + \gamma I)^{\frac{p}{2}})^{\frac{2}{p}}$ ,  $\widehat{\mathcal{W}}^* = \frac{(X^T X + \gamma I)^{\frac{p-1}{2}}}{(\operatorname{Tr}(X^T X + \gamma I)^{\frac{p}{2}})^{\frac{p-2}{p}}}$  satisfy the KKT conditions to (29).

This is because  $\operatorname{Tr}(\widehat{\mathcal{W}}^*)^{\frac{p}{p-2}} = 1$ ,  $\widehat{\mathcal{W}}^* \succ 0$  and  $\lambda^* > 0$ . The complementary slackness is also true since the primal inequality constraint is tight. Since (29) is a convex problem and  $\widehat{\mathcal{W}}^*$  satisfies the KKT conditions, it is the optimal solution to (29). It is also easy to see that  $\tilde{W} = \operatorname{argmin}_{W \succ 0} \mathcal{J}^p(X, W, \gamma)$  where  $\tilde{W} = (X^T X + \gamma I)^{\frac{p}{2}-1}$ . Also note that  $\mathcal{J}^p(X, \tilde{W}, \gamma) = f_p(X)$ . Now,

$$\begin{aligned} \operatorname{argmin}_{Z: \mathcal{A}(Z)=b} \mathcal{J}^p(Z, \tilde{W}, \gamma) &= \operatorname{argmin}_{Z: \mathcal{A}(Z)=b} \operatorname{Tr} \tilde{W} (Z^T Z + \gamma I) \\ &= \operatorname{argmin}_{Z: \mathcal{A}(Z)=b} \operatorname{Tr} (X^T X + \gamma I)^{\frac{p}{2}-1} (Z^T Z + \gamma I) \\ &= \operatorname{argmin}_{Z: \mathcal{A}(Z)=b} \operatorname{Tr} \widehat{\mathcal{W}}^* (Z^T Z + \gamma I) \\ &= \operatorname{argmin}_{Z: \mathcal{A}(Z)=b} \mathcal{F}^p(Z, (\widehat{\mathcal{W}}^*)^{\frac{p}{p-2}}, \gamma). \end{aligned}$$

■

## Appendix B. Proof of Theorem 2

We first present two useful lemmas.

**Lemma 12** For each  $k \geq 1$ , we have

$$\operatorname{Tr}(X^{kT} X^k)^{\frac{p}{2}} \leq \mathcal{J}^p(X^1, W^0, \gamma^0) := D \quad (30)$$

where  $W^0 = I$ ,  $\gamma^0 = 1$ . Also,  $\lambda_j(W^k) \geq D^{(1-\frac{2}{p})}$ ,  $j = 1, 2, \dots, \min\{m, n\}$

**Proof** First, notice that

$$\begin{aligned} \operatorname{Tr}(X^{kT} X^k)^{\frac{p}{2}} &\leq \operatorname{Tr}(X^{kT} X^k + \gamma I)^{p/2} = \mathcal{J}^p(X^k, W^k, \gamma^k) \\ &\leq \mathcal{J}^p(X^1, W^1, \gamma^1) \leq \mathcal{J}^p(X^1, W^0, \gamma^0) = D, \end{aligned}$$

where the second and third inequalities follow from (6). This proves (30). Furthermore, from the above chain of inequalities,

$$(\|X^{kT} X^k\| + \gamma)^{\frac{p}{2}} = \|(X^{kT} X^k + \gamma I)^{\frac{p}{2}}\| \leq D.$$

Using this and the definition of  $W^k$ , we obtain

$$\|(W^k)^{-1}\| = \|(X^{kT} X^k + \gamma I)^{1-\frac{p}{2}}\| = (\|X^{kT} X^k\| + \gamma)^{1-\frac{p}{2}} \leq D^{\left(\frac{2}{p}-1\right)}.$$

This last relation shows that  $\lambda_j(W^k) = \sigma_j(W^k) \geq 1/\|(W^k)^{-1}\| \geq D^{(1-\frac{2}{p})}$  for all  $j$ . ■

**Lemma 13** *A matrix  $X^*$  is a minimizer of*

$$\begin{aligned} & \text{minimize} \quad \text{Tr} W X^T X \\ & \text{subject to} \quad \mathcal{A}(X) = b \end{aligned}$$

*if and only if  $\text{Tr}(W X^{*T} Z) = 0$  for all  $Z \in \mathcal{N}(A)$ .*

**Proof** [of Theorem (2)] For each  $k \geq 1$ , we have that

$$\begin{aligned} 2[\mathcal{J}^p(X^k, W^k, \gamma^k) - \mathcal{J}^p(X^{k+1}, W^{k+1}, \gamma^{k+1})] & \geq 2[\mathcal{J}^1(X^k, W^k, \gamma^k) - \mathcal{J}^1(X^{k+1}, W^k, \gamma^k)] \\ & = \langle X^k, X^k \rangle_{W^k} - \langle X^{k+1}, X^{k+1} \rangle_{W^k} \\ & = \langle X^k + X^{k+1}, X^k - X^{k+1} \rangle_{W^k} \\ & = \langle X^k - X^{k+1}, X^k - X^{k+1} \rangle_{W^k} \\ & = \text{Tr} W^k (X^k - X^{k+1})^T (X^k - X^{k+1}) \\ & \geq D^{(1-\frac{2}{p})} \|X^k - X^{k+1}\|_F^2 \end{aligned}$$

where the above expressions use Lemma 13 and Lemma 12. Summing the above inequalities over all  $k \geq 1$ , we have that  $\lim_{n \rightarrow \infty} (X^n - X^{n+1}) = 0$ . ■

### Appendix C. Proof of Theorem 8

For any two matrices  $X, Y$  we denote  $\langle X, Y \rangle_W = \text{Tr} W X^T Y$ . We first note that the iterates of sIRLS- $p$  satisfy

$$\begin{aligned} \mathcal{J}^p(X^{k+1}, W^{k+1}, \gamma^{k+1}) & \leq \mathcal{J}^p(X^{k+1}, W^k, \gamma^{k+1}) \\ & \leq \mathcal{J}^p(X^{k+1}, W^k, \gamma^k) \\ & \leq \mathcal{J}^p(X^k, W^k, \gamma^k). \end{aligned}$$

The last inequality follows from the Lipschitz continuity (with  $L^k = 2(\gamma^k)^{\frac{p}{2}-1}$ ) of the gradient of  $\text{Tr}(W^k X^T X)$ , that is,

$$\text{Tr} W^k X^T X \leq \text{Tr} W^k X^{kT} X^k + \langle 2X^k W^k, X - X^k \rangle + \frac{L^k}{2} \|X - X^k\|_F^2 \quad \forall X, X^k$$

and the fact that

$$\begin{aligned} X^{k+1} &= \arg \min_X \|X - (X^k - X^k W^k)\|_F^2 \\ &\text{s.t. } \mathcal{P}_\Omega(X) = \mathcal{P}_\Omega(X_0). \end{aligned}$$

The convergence of  $\{\mathcal{J}^p(X^k, W^k, \gamma^k)\}$  follows from Monotone convergence theorem. This also implies that the sequence  $\{X^k\}$  is bounded. Hence there exists a convergent subsequence,  $\{X^{n_i}\} \rightarrow X^*$ . Also let  $\{X^{n_i+1}\} \rightarrow \widehat{X}$ . If  $X^*$  is a stationary point, we are done. Conversely, if  $X^*$  is not a stationary point to (26) then it follows that,  $\widehat{X} \neq X^*$ . But  $\widehat{X} \neq X^*$  implies (using strict convexity) that  $\text{Tr}(W^* \widehat{X}^T \widehat{X}) < \text{Tr}(W^* X^{*T} X^*)$  which also implies that  $\mathcal{J}^p(\widehat{X}, \widehat{W}, \gamma_{\min}) < \mathcal{J}^p(X^*, W^*, \gamma_{\min})$ . However since

$$\begin{aligned} \lim \mathcal{J}^p(X^i, W^i, \gamma^i) &= \lim \mathcal{J}^p(X^{n_i}, W^{n_i}, \gamma^{n_i}) \\ &= \lim \mathcal{J}^p(X^{n_i+1}, W^{n_i+1}, \gamma^{n_i+1}), \end{aligned}$$

we have a contradiction. Therefore,  $X^*$  is a stationary point to (26) and the theorem follows.

## References

- B.J. Dahlen, J.A. Konstan, J. Herlocker, N. Good, A. Borchers, and J. Riedl. “Movie lens data”. 1998. <http://www.grouplens.org/node/73>.
- A. Argyriou. A study of convex regularizers for sparse recovery and feature selection. 2010. Technical Report. Available at [http://ttic.uchicago.edu/~argyriou/papers/sparse\\_report.pdf](http://ttic.uchicago.edu/~argyriou/papers/sparse_report.pdf).
- A. Argyriou, C.A. Micchelli, M. Pontil, and Y. Ying. A spectral regularization framework for multi-task structure learning. In *Proc. of Neural Information Processing Systems (NIPS)*, 2007.
- J.F. Cai, E.J. Candes, and Z. Shen. A singular value thresholding algorithm for matrix completion. *SIAM J. on Optimization*, 20(4):1956–1982, 2008.
- E. J. Candes and S. Becker. Software for singular value thresholding algorithm for matrix completion. 2010. Available at <http://svt.caltech.edu/code.html>.
- E. J. Candes and B. Recht. Exact matrix completion via convex optimization. *Foundations of Computational Mathematics*, 9:717–772, 2009.
- E. J. Candes and T. Tao. Decoding by linear programming. *IEEE Trans Info. Theory*, 2004.
- E. J. Candes, X. Li, Y. Ma, and J. Wright. Robust principal component analysis? *Journal of the ACM*, 58(3), 2011.
- E.J. Candes, M.B. Wakin, and S. Boyd. Enhancing sparsity by reweighted  $l_1$  minimization. *Journal of Fourier Analysis and Applications*, 14:877–905, 2008.

- V. Chandrasekaran, P.A. Parrilo, and A.S. Willsky. Latent variable graphical model selection via convex optimization. *Annals of Statistics*, forthcoming.
- V. Chandrasekaran, B. Recht, P. A. Parrilo, and A. S. Willsky. The convex geometry of linear inverse problems. *Journal of the Foundations of Computational Mathematics*, forthcoming.
- V. Chandrasekaran, S. Sanghavi, P. A. Parrilo, and A. S. Willsky. Rank-sparsity incoherence for matrix decomposition. *SIAM Journal on Optimization*, 21(2):572–596, 2011.
- R. Chartrand and V. Staneva. Restricted isometry properties and nonconvex compressive sensing. *Inverse Problems*, 24(035020):1–14, 2008.
- R. Chartrand and W. Yin. Iteratively reweighted algorithms for compressive sensing. In *33rd International Conference on Acoustics, Speech, and Signal Processing (ICASSP)*, 2008.
- I. Daubechies, R. DeVore, M. Fornasier, and C.S. Gunturk. Iteratively re-weighted least squares minimization for sparse recovery. *Commun. Pure Appl. Math*, 63(1):1–38, 2010.
- P. Drineas, R. Kannan, and M. W. Mahoney. Fast monte carlo algorithms for matrices ii: Computing a low rank approximation to a matrix. *SIAM Journal on Computing*, 36:158–183, 2006.
- M. Fazel, H. Hindi, and S. Boyd. A rank minimization heuristic with application to minimum order system approximation. In *Proc. American Control Conference*, Arlington, VA, 2001.
- M. Fazel, H. Hindi, and S. Boyd. Log-det heuristic for matrix rank minimization with applications to hankel and euclidean distance matrices. In *Proc. American Control Conference*, pages 2156–2162, Denver, CO, 2003.
- M. Fornasier, H. Rauhut, and R. Ward. Low-rank matrix recovery via iteratively reweighted least squares minimization. *SIAM Journal of Optimization*, 21(4), 2011.
- M. Fornasier, H. Rauhut and R. Ward. Code for IRLSM. Available at <http://www.ma.utexas.edu/users/rward/>.
- R. Garg and R. Khandekar. Gradient descent with sparsification: An iterative algorithm for sparse recovery with restricted isometry property. In *Proc. of 26th Intl. Conf. on Machine Learning (ICML)*, 2009.
- D. Goldfarb and S. Ma. Convergence of fixed point continuation algorithms for matrix rank minimization. *Foundations of Computational Mathematics*, 11(2), 2011.
- D. Goldfarb and S. Ma. FPCA code. 2009. Available at <http://www.columbia.edu/~sm2756/FPCA.htm>.
- D. Gross, Y. K. Liu, S. T. Flammia, S. Becker, and J. Eisert. Quantum state tomography via compressed sensing. *Physical Review Letters*, 105, 2010.

- N. Halko, P. G. Martinsson, and J. A. Tropp. Finding structure with randomness: Stochastic algorithms for constructing approximate matrix decompositions. *SIAM Review*, 53(2):217–288, 2011.
- R. A. Horn and C. R. Johnson. *Matrix Analysis*. Cambridge University Press, 1990.
- R. A. Horn and C. R. Johnson. *Topics in Matrix Analysis*. Cambridge University Press, 1991.
- R. H. Keshavan and S. Oh. A gradient descent algorithm on the grassman manifold for matrix completion. 2009. Available at <http://arxiv.org/abs/0910.5260>.
- R. H. Keshavan, A. Montanari, and S. Oh. Optspace algorithm. 2009a. <http://www.stanford.edu/~raghuram/optspace/index.html>.
- R. H. Keshavan, A. Montanari, and S. Oh. Low-rank matrix completion with noisy observations: a quantitative comparison. In *Proc. 47th Annual Allerton Conference on Communication, Control, and Computing*, 2009b.
- K. Lee and Y. Bresler. Admira: Atomic decomposition for minimum rank approximation. *IEEE Tran. Info. Theory*, 56(9), 2010.
- A.S. Lewis. Derivatives of spectral functions. *Mathematics of Operations Research*, 21(3):576–588, 1996.
- Z. Liu and L. Vandenberghe. Interior-point method for nuclear norm approximation with application to system identification. *SIAM J. Matrix Analysis and Appl.*, 31(3), 2008.
- M. S. Lobo, M. Fazel, and S. Boyd. Portfolio optimization with linear and fixed transaction costs. *Annals of Operations Research*, 152:341–365, 2006.
- A. W. Marshall and I. Olkin. *Inequalities: Theory of Majorization and Its Applications*. 1979.
- R. Mazumder, T. Hastie, and R. Tibshirani. Spectral regularization algorithms for learning large incomplete matrices. *Journal of Machine Learning Research*, 11:2287 – 2322, 2010.
- R. Meka, P. Jain, and I. S. Dhillon. Guaranteed rank minimization via singular value projection. In *Proc. of Neural Information Processing Systems (NIPS)*, 2010.
- K. Mohan and M. Fazel. Reweighted nuclear norm minimization with application to system identification. In *Proc. American Control Conference*, Baltimore, MA, 2010a.
- K. Mohan and M. Fazel. Iterative reweighted least squares for matrix rank minimization. In *Proc. 48th Allerton Conference on Controls and Communications*, Allerton, IL, 2010b.
- D. Needell. Noisy signal recovery via iterative reweighted  $\ell_1$ -minimization. In *Proc. Asilomar Conference on Signals, Systems and Computers*, 2009.



- D. Needell and J. A. Tropp. Cosamp: Iterative signal recovery from incomplete and inaccurate samples. *Applied and Computational Harmonic Analysis*, 26(3):301–321, 2008.
- S. Oymak and B. Hassibi. New null space results and recovery thresholds for matrix rank minimization. 2010. Available at <http://arxiv.org/abs/1011.6326>.
- S. Oymak, K. Mohan, M. Fazel, and B. Hassibi. A simplified approach to recovery conditions for low rank matrices. In *Proc. International Symposium on Information Theory*, 2011.
- B. D. Rao and K. Kreutz-Delgado. An affine scaling methodology for best basis selection. *IEEE Transactions on Signal Processing*, 47:187–200, 1999.
- N. Srebro, J.D.M. Rennie, and T.S. Jakkola. Maximum-margin matrix factorization. In *Advances in Neural Information Processing Systems*, 2005b.
- W. Tan, G. Cheung, and Y. Ma. Face recovery in conference video streaming using robust principal component analysis. In *Proc. IEEE Intl. Conf. on Image Processing*, 2011.
- K. C. Toh and S. W. Yun. An accelerated proximal gradient algorithm for nuclear norm regularized least squares problem. *Pacific Journal of Optimization*, 6:615–640, 2010.
- D. P. Wipf and S. Nagarajan. Iterative reweighted  $\ell_1$  and  $\ell_2$  methods for finding sparse solutions. *Journal of Selected Topics in Signal Processing (Special Issue on Compressive Sensing)*, 4(2), 2010.
- T. Zhang. Analysis of multi-stage convex relaxation for sparse regularization. *Journal of Machine Learning Research*, 11:1081–1107, 2010.